

Modern quantum many-body physics – Interacting bosons

Xiao-Gang Wen

The first step to build a theory: how to label states?

One particle states

- How to label states of one boson in 1D space? $\rightarrow |x\rangle$. The most general state $|\psi\rangle = \int dx \psi(x) |x\rangle$
- Energy eigenstates (momentum eigenstates) $|k\rangle = \int dx e^{ikx} |x\rangle$, where wave vector $k = \text{int.} \times \frac{2\pi}{L}$. (The space is a 1D ring of size L)
 - Momentum = $p = \hbar k$.
 - Energy = $\epsilon_k = \frac{\hbar^2 k^2}{2M}$ (Or $\epsilon_k = \hbar |k|c$ for massless photons)

Many-particle states

- Label all zero-, one-, two-, three-, ... boson states:

$$|\emptyset\rangle$$

$$|k_1\rangle$$

$$|k_1, k_2\rangle, k_1 \leq k_2 \quad (|k_1, k_2\rangle = |k_2, k_1\rangle \text{ for identical particles})$$

$$|k_1, k_2, k_3\rangle, k_1 \leq k_2 \leq k_3$$

... ..

- Label all zero-, one-, two-, three-, ... boson states

(The **second quantization** – quantum field theory of bosons):

$n_k \equiv$ the number of bosons with wave vector k .

$|\{n_k = 0\}\rangle$ is the ground state. $|\{n_k \neq 0\}\rangle$ is an excited state.

$|\{n_k = 0\}\rangle = |\emptyset\rangle$. No boson

$|\{n_{k_1} = 1, \text{others} = 0\}\rangle = |k_1\rangle$. One boson





$|\{n_{k_1} = 1, n_{k_2} = 1, \text{others} = 0\}\rangle = |k_1, k_2\rangle = |k_2, k_1\rangle$.

$|\{n_{k_1} = 1, n_{k_2} = 1, n_{k_3} = 1, \text{others} = 0\}\rangle = |k_1, k_2, k_3\rangle = |k_2, k_3, k_1\rangle = \dots$

$|\{n_{k_1} = 2, n_{k_2} = 1, \text{others} = 0\}\rangle = |k_1, k_1, k_2\rangle = |k_1, k_2, k_1\rangle = \dots$

A many-boson system with no interaction = a collection of decoupled harmonic oscillators

$n_k \rightarrow$ the occupation number of the bosons on orbital- k .



- If we ignore the interaction between bosons $|\{n_k\}\rangle$ is an energy eigenstate with energy $E = \sum_k n_k \epsilon_k$
- The above energy can be viewed as the total energy of a collection of decoupled harmonic oscillators. The oscillators are labeled by $k = \text{int.} \times \frac{2\pi}{L}$. The oscillator labeled by k has a frequency $\omega_k = \epsilon_k/\hbar$.
- A collection of decoupled harmonic oscillators = vibration modes of a vibrating string. The two polarizations of bosons \rightarrow two directions of string vibrations
 \rightarrow **quantum field theory** of 1D boson gas.

Many-body Hamiltonian for non-interacting bosons

View 1D non-interacting bosons (with $0, 1, 2, 3, \dots$ bosons) as a collection of oscillators with frequencies ω_k :

$$\hat{H} = \sum_k \left(\hat{a}_k^\dagger \hat{a}_k + \frac{1}{2} \right) \hbar \omega_k, \quad \hbar \omega_k = \epsilon_k = \frac{\hbar^2 k^2}{2m}, \quad k = \text{int.} \times \frac{2\pi}{L}$$

raising-lowering operator

$$\hat{a}_k = \sqrt{\frac{m\omega_k}{2\hbar}} \left(\hat{x}_k + \frac{i}{m\omega_k} \hat{p}_k \right), \quad [\hat{a}_k, \hat{a}_{k'}^\dagger] = \delta_{k,k'}$$

$$\hat{a}_k^\dagger \hat{a}_k |n_k\rangle = n_k |n_k\rangle, \quad \hat{a}_k^\dagger |n_k\rangle = |n_k + 1\rangle, \quad \hat{a}_k |n_k\rangle = |n_k - 1\rangle.$$

- All the energy eigenstates are labeled by $|\{n_k\}\rangle = \bigotimes_k |n_k\rangle$.
The total energy $E_{\text{tot}} = \sum_k (n_k + \frac{1}{2}) \epsilon_k$.
The total particle number $N = \sum_k n_k$.

$\hat{a}_k^\dagger, \hat{a}_k$ are also creation-annihilation operator of bosons.

Many-body Hamiltonian for bosons on lattice

- Infinite problem on quantum field theory:
The vacuum energy $E_0 = 0$ or $E_0 = \sum_k \frac{1}{2}\epsilon_k$?
The right answer $E_0 = \sum_k \frac{1}{2}\epsilon_k = \infty$
- Non-interacting bosons on a lattice

For 1D non-interacting bosons
(with 0, 1, 2, 3, ... bosons)


$$\hat{H} = \sum_{k \in BZ} \left(\hat{a}_k^\dagger \hat{a}_k + \frac{1}{2} \right) \epsilon_k, \quad \epsilon_k = 2t[1 - \cos(ka)],$$

$$k = \text{int.} \times \frac{2\pi}{L} \in \left[-\frac{\pi}{a}, \frac{\pi}{a}\right].$$

- The vacuum energy now is finite

$$E_0 = \sum_{k \in BZ} \frac{1}{2} \epsilon_k = L \int_{-\frac{\pi}{a}}^{\frac{\pi}{a}} \frac{dk}{2\pi} 2t[1 - \cos(ka)] = L \frac{2t}{a} = 2tN.$$

- The vacuum energy can be measured via **Casimir effect**.



© Source unknown. All rights reserved. This content is excluded from our Creative Commons license. For more information, see <https://ocw.mit.edu/help/faq-fair-use>.

Many-body Hamiltonian for interacting bosons on lattice

- The total particle number operator

$$\hat{N} = \sum_{k \in \text{BZ}} \hat{a}_k^\dagger \hat{a}_k = \sum_i \hat{\varphi}_i^\dagger \hat{\varphi}_i, \quad [\hat{\varphi}_i, \hat{\varphi}_j^\dagger] = \delta_{ij}.$$

$$\hat{a}_k = \sum_{x_i} N^{-1/2} e^{ikx_i} \hat{\varphi}_i, \quad x_i = a i, \quad i = 1, \dots, N;$$

- $\hat{n}_k = \hat{a}_k^\dagger \hat{a}_k$ is the number operator for bosons on orbital k .
- $\hat{n}_i = \hat{\varphi}_i^\dagger \hat{\varphi}_i$ is the number operator for bosons on site i . $\hat{\varphi}_i^\dagger, \hat{\varphi}_i$ are creation-annihilation operator of bosons at site- i .

- Many-body Hamiltonian for interacting bosons

$$H = \sum_k (\hat{a}_k^\dagger \hat{a}_k + \frac{1}{2}) \epsilon_k - \sum_i \mu \hat{n}_i + \sum_{i < j} V_{ij} \hat{n}_i \hat{n}_j$$

$$= \sum_k \frac{1}{2} (\hat{a}_k^\dagger \hat{a}_k + \hat{a}_k \hat{a}_k^\dagger) \epsilon_k - \sum_i \mu \hat{\varphi}_i^\dagger \hat{\varphi}_i + \sum_{i < j} V_{ij} \hat{\varphi}_i^\dagger \hat{\varphi}_i \hat{\varphi}_j^\dagger \hat{\varphi}_j$$

$$= \sum_i \left[t (\hat{\varphi}_i^\dagger \hat{\varphi}_i + \hat{\varphi}_i \hat{\varphi}_i^\dagger) - t (\hat{\varphi}_{i+1}^\dagger \hat{\varphi}_i + \hat{\varphi}_i^\dagger \hat{\varphi}_{i+1}) \right] - \sum_i \mu \hat{\varphi}_i^\dagger \hat{\varphi}_i + \sum_{i < j} V_{ij} \hat{\varphi}_i^\dagger \hat{\varphi}_i \hat{\varphi}_j^\dagger \hat{\varphi}_j$$

Hard-core bosons and spin-1/2 systems

- Assume on-site interaction $V_{ij} = U\delta_{ij}$, $\mu = U + 2B + t \rightarrow U\hat{n}_i\hat{n}_i - \mu\hat{n}_i = U(\hat{n}_i - 1)\hat{n}_i - (2B + t)\hat{n}_i$, $U \rightarrow +\infty$

The low energy sector for interaction $\rightarrow n_i = 0, 1$ (\downarrow, \uparrow) or

$$n_i = \frac{\sigma_i^z - 1}{2}, \quad \hat{\varphi}_i = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} = \sigma_i^- = \frac{\sigma_i^x - i\sigma_i^y}{2}.$$

Hamiltonian for interacting bosons = a spin-1/2 system

$$\begin{aligned} H_{XY\text{-model}} &= \sum_i \left[-t(\sigma_i^+ \sigma_{i+1}^- + \sigma_i^- \sigma_{i+1}^+) - B\sigma_i^z \right] \\ &= \sum_i \left[-J(\sigma_i^x \sigma_{i+1}^x + \sigma_i^y \sigma_{i+1}^y) - B\sigma_i^z \right], \quad J = \frac{1}{2}t \end{aligned}$$

- $U(1)$ symmetry generated by $U_\phi = \prod_i e^{i\phi\sigma_i^z/2}$: $U_\phi H U_\phi = H$.
 $\sum_i \sigma_i^z \sim N + \text{const.}$ conservation.
- Phase diagram: Treat operators σ as classical unit-vector (spin) \mathbf{n} .

$$B < 0 : |\downarrow \cdots \downarrow\rangle \quad B \sim 0 : |\rightarrow \cdots \rightarrow\rangle \quad B > 0 : |\uparrow \cdots \uparrow\rangle$$

————— **—————** ————— $\rightarrow B$

0-boson/site

Superfluid

1-boson/site (Mott insulator)

Hard-core bosons and spin-1 systems

- Assume on-site interaction to have a form $U[(n_i - 1)^4 - (n_i - 1)^2]$.
The low energy sector for the interaction: $n_i = 0, 1, 2$ ($\downarrow, 0, \uparrow$) or

$$n_i = S_i^z - 1, \quad \hat{\phi}_i = S_i^-.$$

Hamiltonian for interacting bosons = a spin-1 system ($U(1)$ symm.)

$$\begin{aligned} H &= \sum_i \left[-t(S_i^+ S_{i+1}^- + S_i^- S_{i+1}^+) - BS_i^z + V(S_i^z)^2 \right] \\ &= \sum_i \left[-J(S_i^x S_{i+1}^x + S_i^y S_{i+1}^y) - BS_i^z + V(S_i^z)^2 \right]. \end{aligned}$$

- B - V phase diagram Treat operators σ as classical unit-vector (spin) \mathbf{n} .


- Two different critical points:

- The black-line represents a $z = 2$ critical point.
(ie excitations have dispersion relation $\omega_k \sim k^2$)

- The filled dot represents a different

$z = 1$ critical point with emergent Lorentz symmetry

(ie excitations have dispersion relation $\omega_k \sim k$)



$J = 1$

Many-body Hamiltonian

- Consider a system formed by two spin-1/2 spins. The spin-spin interaction: $H = J(\sigma_1^x \sigma_2^x + \sigma_1^y \sigma_2^y + \sigma_1^z \sigma_2^z)$.

where $\sigma_i^{x,y,z}$ are the Pauli matrices acting on the i^{th} spin.

$J < 0 \rightarrow$ ferromagnetic, $J > 0 \rightarrow$ antiferromagnetic.

Is H a two-by-two matrix? In fact

$$H = -J[(\sigma^x \otimes I) \cdot (I \otimes \sigma^x) + (\sigma^y \otimes I) \cdot (I \otimes \sigma^y) + (\sigma^z \otimes I) \cdot (I \otimes \sigma^z)]$$

H is a four-by-four matrix:

$$\sigma_1^z \sigma_2^z = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \quad \sigma_1^x \sigma_2^x = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix} \quad \sigma_1^x \sigma_2^y = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \\ 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{pmatrix}$$

- $\sigma_i^z = I \otimes \dots \otimes I \otimes \sigma^z \otimes I \otimes \dots \otimes I$ is a $2^{N_{\text{site}}}$ -dimensional matrix

Example: An 1D ring formed by L spin-1/2 spins:

$$H = - \sum_{i=1}^L \sigma_i^x \sigma_{i+1}^x - h \sum_{i=1}^L \sigma_i^z$$

– transverse Ising model. H is a $2^L \times 2^L$ matrix.

Condensed matter: A local many-body quantum system

- A many-body quantum system
= Hilbert space \mathcal{V}_{tot} + Hamiltonian H

- The locality of the Hilbert space:

$$\mathcal{V}_{tot} = \otimes_{i=1}^N \mathcal{V}_i$$



- The i also label the vertices of a graph

- A local Hamiltonian $H = \sum_{\mathbf{x}} H_{\mathbf{x}}$ and $H_{\mathbf{x}}$ are local hermitian operators acting on a few neighboring \mathcal{V}_i 's.

- A quantum state, a vector in \mathcal{V}_{tot} :

$$|\Psi\rangle = \sum \Psi(m_1, \dots, m_N) |m_1\rangle \otimes \dots \otimes |m_N\rangle, \\ |m_i\rangle \in \mathcal{V}_i$$

- A gapped Hamiltonian has the following spectrum as $N \rightarrow \infty$
(eg $H = -\sum (J\sigma_i^z \sigma_{i+\delta}^z + h\sigma_i^x)$)



Many-body spectrum using Octave (Matlab or Julia)


Transverse Ising model on a ring of L site:

$$H = -J \sum_{i=1}^L \sigma_i^x \sigma_{i+1}^x - h \sum_{i=1}^L \sigma_i^z$$

H is an 2^L -by- 2^L matrix, whose eigenvalues can be computed via the following Octave code (the code also run in Matlab or Julia with minor modifications):

```
X=sparse([0,1;1,0]); Z=sparse([1,0;0,-1]); XX=kron(X,X);
L=13; h=1.0; J=1.0
H=-kron(kron(X, speye(2^(L-2)) ),X);
for i=1:L-1
    H=H - kron( speye(2^(i-1)), kron(J*XX, speye(2^(L-1-i))) );
end
for i=1:L
    H=H - kron( speye(2^(i-1)), kron(h*Z, speye(2^(L-i))) );
end
eigs( H , 10, 'sa') # compute the lowest 10 eigenvalues
```

The 100 lowest energy eigenvalues for $L = 16$, as functions of $h/J \in [0, 2]$.



Quantum phases and quantum phase transitions


- Phases are defined through phase transitions.

What are phase transitions?


As we change a parameter g in Hamiltonian $H(g)$, the ground state energy density $\epsilon_g = E_g/V$ or the average of a local operator $\langle \hat{O} \rangle$ may have a singularity at g_c : the system has a phase transition at g_c .

The Hamiltonian $H(g)$ is a smooth function of g . How can the ground state energy density ϵ_g be singular at a certain g_c ?

- There is no singularity for finite systems. Singularity appears only for infinite systems.
- Spontaneous symmetry breaking is a mechanism to cause a singularity in ground state energy density ϵ_g .
→ Spontaneous symmetry breaking causes phase transition.



$E_2 - E_1$ of trans. Ising
for $L = 3, \dots, 13$



Symmetry breaking theory of phase transition


It is easier to see a phase transition in the semi classical approximation of a quantum theory.

- Variational ground state $|\Psi_\phi\rangle$ for H_g is obtained by minimizing energy $\epsilon_g(\phi) = \frac{\langle \Psi_\phi | H_g | \Psi_\phi \rangle}{V}$ against the variational parameter ϕ .
 $\epsilon_g(\phi)$ is a smooth function of ϕ and g . How can its minimal value $\epsilon_g \equiv \epsilon_g(\phi_{min})$ have singularity as a function of g ?

- Minimum splitting \rightarrow singularity in $\frac{\partial^2 \epsilon_g}{\partial g^2}$ at g_c . Second order trans.
State-B has less symmetry than state-A.
State-A \rightarrow State-B: spontaneous symmetry breaking.

- For a long time, we believe that

phase transition = change of symmetry
the different phases = different symmetry.



- Minimum switching \rightarrow singularity in $\frac{\partial \epsilon_g}{\partial g}$ at g_c . First order trans.


Example: meanfield symmetry breaking transition

Consider a transverse field Ising model $H = \sum_i -J\sigma_i^x\sigma_{i+1}^x - h\sigma_i^z$

Use trial wave function $|\Psi_\phi\rangle = \otimes_i |\psi_i\rangle$, $|\psi_i\rangle = \cos \frac{\phi}{2} |\uparrow\rangle + \sin \frac{\phi}{2} |\downarrow\rangle$ to estimate the ground state energy

$$\begin{aligned}\langle \Psi_\phi | H | \Psi_\phi \rangle &= - \sum \langle \psi_i | \sigma_i^x | \psi_i \rangle \langle \psi_{i+1} | \sigma_{i+1}^x | \psi_{i+1} \rangle - h \sum \langle \psi_i | \sigma_i^z | \psi_i \rangle \\ &= (2J \cos \frac{\phi}{2} \sin \frac{\phi}{2})^2 - h(\cos^2 \frac{\phi}{2} - \sin^2 \frac{\phi}{2}) = \sin^2 \phi - h \cos \phi\end{aligned}$$

Phase transition at $h/J = 2$. ($h/J = 1.5, 2.0, 2.5$)



Order parameter and symmetry-breaking phase transition

ϕ or σ_i^x are order parameters for the Z_2 symm.-breaking transition:

- Under Z_2 ($180^\circ S^z$ rotation), $\phi \rightarrow -\phi$ or $\sigma_i^x \rightarrow -\sigma_i^x$
- In symmetry breaking phase $\phi = \pm\phi_0$, $\langle \sigma_i^x \rangle = \pm$.

In symmetric phase $\phi = 0$, $\langle \sigma_i^x \rangle = 0$. (**Classical picture**)

Ginzberg-Landau theory of continuous phase transition

- Quantum Z_2 -Symmetry: generator $U = \prod_j \sigma_j^z$, $U^2 = 1$.
Symmetry trans.: $U\sigma_i^z U^\dagger = \sigma_i^z$, $U\sigma_i^x U^\dagger = -\sigma_i^x$, $U\sigma_i^y U^\dagger = -\sigma_i^y$.
 $\rightarrow UHU^\dagger = H$. If $H|\psi\rangle = E_{\text{grnd}}|\psi\rangle$, then $UH|\psi\rangle = E_{\text{grnd}}U|\psi\rangle \rightarrow UHU^\dagger U|\psi\rangle = E_{\text{grnd}}U|\psi\rangle \rightarrow HU|\psi\rangle = E_{\text{grnd}}U|\psi\rangle$
Both $|\psi\rangle$ and $U|\psi\rangle$ are ground states of H :
Either $|\psi\rangle \propto U|\psi\rangle$ (symmetric) or $|\psi\rangle \not\propto U|\psi\rangle$ (symm.-breaking).
- Trial wave function $|\Psi_\phi\rangle = \bigotimes_i (\cos \frac{\phi}{2} |\uparrow\rangle_i + \sin \frac{\phi}{2} |\downarrow\rangle_i)$: $U|\Psi_\phi\rangle = |\Psi_{-\phi}\rangle$
 $\rightarrow \langle \Psi_\phi | H | \Psi_\phi \rangle = \langle \Psi_\phi | U^\dagger U H U^\dagger U | \Psi_\phi \rangle = \langle \Psi_{-\phi} | H | \Psi_{-\phi} \rangle \rightarrow \epsilon(h, \phi) = \epsilon(h, -\phi)$
- If $|\Psi_{\phi=0}\rangle$ is the ground state \rightarrow symmetric phase.
If $|\Psi_{\phi \neq 0}\rangle$ is the ground state \rightarrow symmetry breaking phase.
- Near the phase transition ϕ is small \rightarrow

$$\epsilon(h, \phi) = \epsilon_0(h) + \frac{1}{2}a(h)\phi^2 + \frac{1}{4}b(h)\phi^4 + \dots$$

Transition happen at $a(h_c) = 0$.

Properties near the $T = 0$ (quantum) phase transition

- Ground state energy density:

$$\phi = 0, \quad \epsilon_{\text{grnd}}(h) = \epsilon_0(h) \text{ if } a(h) > 0$$

$$\phi = \pm \sqrt{\frac{-a}{b}}, \quad \epsilon_{\text{grnd}}(h) = \epsilon_0(h) - \frac{1}{4} \frac{a(h)^2}{b} \text{ if } a(h) < 0$$

$\epsilon_{\text{grnd}}(h)$ is non-analytic at the transition point: $a(h) = a_0(h - h_c)$:

$$\epsilon_{\text{grnd}}(h) = \begin{cases} \epsilon_0(h), & h > h_c \\ \epsilon_0(h) - \frac{1}{4} \frac{a_0(h-h_c)^2}{b}, & h < h_c \end{cases}$$

- Magnetization in z -direction: $M_z = \frac{\partial \epsilon_{\text{grnd}}(h)}{\partial h}$.

$$M_z = \frac{\partial \epsilon_0(h)}{\partial h}, \quad h > h_c$$

$$M_z = \frac{\partial \epsilon_0(h)}{\partial h} - \frac{1}{2} \frac{a_0(h-h_c)}{b}, \quad h < h_c$$

$$\rightarrow \Delta M_z \sim |\Delta h|$$


- Magnetization in x -dir.: $M_x = \langle \sigma^x \rangle = \sin \phi$

$$\phi = \pm \sqrt{\frac{-a(h)}{b}} \rightarrow \Delta M_x \sim |\Delta h|^{1/2}$$

- Magnetic susceptibility in x -direction:

$$\text{From } \epsilon(h, \phi, h_x) = \frac{1}{2} a(h) \phi^2 - h_x \phi + \dots$$

$$\rightarrow M_x = \phi = \frac{1}{a(h)} \rightarrow \chi_x = \frac{1}{a(h)} \rightarrow \Delta \chi_x \sim |\Delta h|^{-1}$$



Quantum picture of continuous phase transition

No symmetry breaking in quantum theory according: If $[H, U] = 0$, then H and U share a common set of eigenstates. The ground state $|\Psi_{\text{grnd}}\rangle$ of H , is an eigenstate of U : $U|\Psi_{\text{grnd}}\rangle = e^{i\theta}|\Psi_{\text{grnd}}\rangle$.

No symmetry breaking.

$|\Psi_{\phi}\rangle$ and $|\Psi_{-\phi}\rangle$ in semi classical approximation are not true ground states. The true ground state is $|\Psi_{\text{grnd}}\rangle = |\Psi_{\phi}\rangle + |\Psi_{-\phi}\rangle$ which do not break the symmetry.

- **Quantum picture:** Symmetry-breaking order parameter is zero, $\langle \Psi_{\text{grnd}} | \sigma_i^x | \Psi_{\text{grnd}} \rangle = 0$, for the true ground state. But **the ground states**, $|\Psi_{\text{grnd}}\rangle = |\Psi_{\phi}\rangle + |\Psi_{-\phi}\rangle$ and $|\Psi'_{\text{grnd}}\rangle = |\Psi_{\phi}\rangle - |\Psi_{-\phi}\rangle$, **have an exponentially small energy separation** $\Delta \sim e^{-L/\xi}$.

Symmetry-breaking order parameter is non-zero only for approximate ground states, $|\Psi_{\phi}\rangle$ and $|\Psi_{-\phi}\rangle$.

- **Detect symmetry breaking from correlation function:**

$$\lim_{|i-j| \rightarrow \infty} \langle \Psi_{\text{grnd}} | \sigma_i^x \sigma_j^x | \Psi_{\text{grnd}} \rangle = \text{const.}$$

$$\text{Symmetric phase: } \lim_{|i-j| \rightarrow \infty} \langle \Psi_{\text{grnd}} | \sigma_i^x \sigma_j^x | \Psi_{\text{grnd}} \rangle = 0$$

Collective mode of order parameter ϕ : guess

- From the energy $\epsilon(h, \phi) = \epsilon_0(h) + \frac{1}{2}a(h)\phi^2 + \frac{1}{4}b(h)\phi^4 + \dots$
 \rightarrow Restoring force $f = -a\phi - b\phi^3 \rightarrow$ EOM $\rho\ddot{\phi} = -a\phi - b\phi^3$.
- $k \neq 0$ mode: $\epsilon(h, \phi) = \frac{1}{2}g(\partial_x\phi)^2 + \frac{1}{2}a(h)\phi^2 + \frac{1}{4}b(h)\phi^4 + \dots$
 Restoring force $f = g\partial_x^2\phi - a\phi - b\phi^3$
 \rightarrow EOM $\rho\ddot{\phi} = g\partial_x^2\phi - a\phi - b\phi^3$.


- Collective mode: $\omega_k = \sqrt{\frac{gk^2+a}{\rho}}$
 Energy gap: $\Delta = \sqrt{\frac{a(h)}{\rho}} = \sqrt{\frac{a_0(h-h_c)}{\rho}}$.

- At the critical point $h = h_c$:
 Gapless = diverging susceptibility

$\omega_k \sim k^z$, $z = 1$. z is the **dynamical critical exponent**.

$z = 1 \rightarrow$ Emergence of Lorentz symmetry.

Where does ρ come from?



Continuous quantum phase transition between gapped phases = gap closing phase transition. Continuous quantum phase transition between gapless phases : more low energy modes at the critical point.

Collective mode of order parameter ϕ : calculate

Consider a transverse field Ising model $H = -\sum_i (J\sigma_i^x\sigma_{i+1}^x + h\sigma_i^z)$.

Trial wave function $|\Psi_{\phi_i}\rangle = \otimes_i |\phi_i\rangle$, $|\phi_i\rangle = \frac{|\uparrow\rangle + \phi_i|\downarrow\rangle}{\sqrt{1+|\phi_i|^2}}$ (Key: ϕ_i complex)

$$\langle\sigma_i^x\rangle = \frac{\phi_i + \phi_i^*}{1 + |\phi_i|^2}, \quad \langle\sigma_i^z\rangle = \frac{1 - |\phi_i|^2}{1 + |\phi_i|^2}.$$

- Average energy

$$\bar{H} = -\sum_i \left[J \frac{(\phi_i + \phi_i^*)(\phi_{i+1} + \phi_{i+1}^*)}{(1 + |\phi_i|^2)(1 + |\phi_{i+1}|^2)} + h \frac{1 - |\phi_i|^2}{1 + |\phi_i|^2} \right]$$

Geometric phase term

$$\begin{aligned} \langle\phi_i|\frac{d}{dt}|\phi_i\rangle &= \frac{\dot{\phi}_i^* \phi_i}{1 + |\phi_i|^2} + (1 + |\phi_i|^2)^{1/2} \frac{d}{dt} (1 + |\phi_i|^2)^{-1/2} \\ &= \frac{\dot{\phi}_i^* \phi_i}{1 + |\phi_i|^2} - \frac{1}{2} \frac{d}{dt} \log(1 + |\phi_i|^2) \end{aligned}$$

Phase space Lagrangian (quadratic approximation: $\phi_i = q_i + \frac{i}{2}p_i$ small)

$$\begin{aligned} L &= \langle\Phi_{\phi_i}|\dot{i} \frac{d}{dt} - H|\Phi_{\phi_i}\rangle = \sum_i i\dot{\phi}_i^* \phi_i + J(\phi_i + \phi_i^*)(\phi_{i+1} + \phi_{i+1}^*) - 2h|\phi_i|^2 \\ &= \sum_i \left[p_i \dot{q}_i + 4Jq_i q_{i+1} - 2h(q_i^2 + \frac{1}{4}p_i^2) \right] \end{aligned}$$

Collective mode of order parameter ϕ : calculate

EOM:

$$\dot{q}_i = \frac{\partial \bar{H}}{\partial p_i} = \frac{\hbar}{2} p_i, \quad \dot{p}_i = -\frac{\partial \bar{H}}{\partial q_i} = 4J(q_{i+1} + q_{i-1}) - 4\hbar q_i$$

in k -space ($q_i = \sum_k N^{-1/2} e^{ikia} q_k$, $p_i = \sum_k N^{-1/2} e^{ikia} p_k$):

$$\dot{q}_k = \frac{\hbar}{2} p_k, \quad \dot{p}_k = 4(Je^{ika} + Je^{-ika} - \hbar)q_k$$

k label harmonic oscillators with EOM

$$\ddot{q}_k = 2\hbar[2\cos(ka) - \hbar]q_k \rightarrow -\omega_k^2 = 2\hbar[2J\cos(ka) - \hbar]$$

The dispersion of the collective mode

$$\omega_k = \sqrt{2\hbar[h - 2J\cos(ka)]}$$

- For $h > 2J$, gap = $\sqrt{2\hbar(h - 2J)}$.

For $h = 2J$, gapless mode with velocity $v = 2aJ$ and $\omega_k = v|k|$.

Many-body spectrum at the critical point

- At the critical point, the gapless excitation is described by a real scalar field ϕ (or q_i) with EOM:


$$\ddot{\phi} = v^2 \partial_x^2 \phi.$$

= an oscillator for every $k = \frac{2\pi}{L} n$

= a wave mode with $\omega_k = v|k|$

= a boson with $\epsilon(p) = v|p|$

- Many-body spectrum for right movers:




Do not count for the $k=0$ orbital.

- Total energy and total momentum for right movers $E = vK$.

Magic at critical point: Emergence of **Lorentz invariance** $\epsilon = vk$.

Emergence of independent right-moving and left-moving sectors (extra degeneracy in many-body spectrum): **conformal invariance**



$z = 1$ and $z = 2$ critical points


The transverse Ising model, $H = -\sum_i (J\sigma_i^x \sigma_{i+1}^x + h\sigma_i^z)$,

has $z = 1$ critical points at $h = \pm J$

The spin-1 XY model,

$H = \sum_i (-JS_i^x S_{i+1}^x - JS_i^y S_{i+1}^y + V(S_i^z)^2 - BS_i^z)$,

has $z = 1$ and $z = 2$ critical points.



- The $z = 1$ critical point appears when $B = 0$ and the spin-1 XY model has the $S^z \rightarrow -S^z$ symmetry.
- The phase space Lagrangian of has a form $\mathcal{L} = A\dot{\phi}^* \dot{\phi} + B\dot{\phi}^* \phi - C|\partial\phi|^2$ for the collective mode at the critical point. When $B = 0$, $A = 0$, which leads to the $z = 1$ critical point. When $B \neq 0$, $A \neq 0$, which leads to the $z = 2$ critical point.

The minimal value of dynamical exponent z is 1

- The $z = 2$ critical point can appear if we have $U(1)$ spin rotation symmetry in the S^x - S^y plane. In this case, the critical point describe the transition from a gapped Mott insulator (spin polarized) phase to a gapless superfluid (XY spin order) phase ($U(1)$ symmetry breaking phase) with $z = 1$ (ie $\omega \sim k$).
- The gapless is the Goldstone mode. **Spontaneous breaking of a continuous symmetry always give rise to a gapless model.**
- The critical point always has more low energy excitations then the two phases it connects.
- The $z = 1$ critical point can appear if we have Z_2 spin rotation symmetry in the $S^x \rightarrow -S^x$. In this case, the critical point describe the transition from a gapped symmetric phase to a gapped spontaneous Z_2 -symmetry breaking phase.
- $z < 1$, $\omega \sim |k|^z$ **is not allowed for short range interaction**, since the velocity for any excitations has an upper bound $v \lesssim a \|H_{i,i+a}\|/\hbar$

The property of $k = 0$ mode (quadratic approx. valid?)

- Now consider transverse Ising model in d dimensions ($g \sim J, h$)

$$L = \sum_i \sum_{\mu=x,y,\dots} [p_i \dot{q}_i + 4Jq_i q_{i+\mu}] - \sum_i [2h(q_i^2 + \frac{1}{4}p_i^2) - gq_i^4]$$

The transition point now is at $h = 2dJ$

- At the critical point $h = 2dJ$,
the $k = 0$ mode is described by the Lagrangian

$$\begin{aligned} L &= Np\dot{q} - \frac{N}{2}hp^2 - Ngq^4 \\ &= \tilde{p}\dot{\tilde{q}} - \frac{h}{2}\tilde{p}^2 - \frac{g}{N}\tilde{q}^4, \quad \tilde{p} = \sqrt{N}p, \quad \tilde{q} = \sqrt{N}q. \end{aligned}$$

- The zero-point energy from the $k = 0$ mode $\tilde{p}\tilde{q} \sim 1 \rightarrow \tilde{q} \sim N^{1/6}$
minimizing: $\frac{h}{2}\tilde{p}^2 + \frac{g}{N}\tilde{q}^4 \sim \frac{h}{2}\tilde{q}^{-2} + \frac{g}{N}\tilde{q}^4 \sim JN^{-1/3}$

The non-linear term is important for $k = 0$ mode.

- The zero-point energy from the k mode (ignoring the non-linear term)
 $Jk \sim JN^{-1/d} |_{k \sim N^{-1/d}}$

The non-linear effect for k mode

- At the critical point $h = 2dJ$, the k mode is described by the Lagrangian

$$\begin{aligned} L &= Np\dot{q} - JNk^2q^2 - \frac{N}{2}hp^2 - Ngq^4 \\ &= \tilde{p}\dot{\tilde{q}} - Jk^2\tilde{q}^2 - \frac{h}{2}\tilde{p}^2 - \frac{g}{N}\tilde{q}^4, \quad \tilde{p} = \sqrt{N}p, \quad \tilde{q} = \sqrt{N}q. \end{aligned}$$

- The zero-point energy from the k mode $\tilde{p}\tilde{q} \sim 1 \rightarrow \tilde{p} \sim 1/\tilde{q} \sim \sqrt{k}$

$$Jk^2\tilde{q}^2 + \frac{h}{2}\tilde{p}^2 + \frac{g}{N}\tilde{q}^4 \sim Jk + \frac{h}{2}k + \frac{g}{Nk^2}$$

The non-linear term is important if

$$\frac{g}{Nk^2} > Jk \quad \text{or} \quad k < \frac{1}{N^{1/3}}$$

- Since the smallest k is $\frac{1}{N^{1/d}}$. For $d > 3$ there is no k satisfying the above condition (except $k = 0$). We can ignore the non-linear term.

Our critical theory from quadratic approximation is correct.

- For $d \leq 3$, we cannot ignore the non-linear term.

Our critical theory from quadratic approximation is incorrect.

Quantum fluctuations: relevant/irrelevant perturbations

EOM of Z_2 order parameter for the $d + 1$ D-transverse Ising model

$$\rho \ddot{\phi} = g \partial_x^2 \phi + a \phi + b \phi^3$$

Is the $b\phi^3$ term important at the transition point $a = 0$?

- The action $S = \int dt d^d x [\frac{1}{2} \rho (\dot{\phi})^2 - \frac{1}{2} g (\partial_x \phi)^2 - \frac{1}{2} a \phi^2 - \frac{1}{4} b \phi^4]$
 - Treating the above as a quantum system with quantum fluctuations, the term $\frac{1}{4} b \phi^4$ is irrelevant if dropping it does not affect the low energy properties at critical point $a = 0$. Otherwise, it is relevant.
 - Rescale t to make $\rho = g$ and rescale ϕ to make $\rho = g = 1$.
 - Consider the fluctuation at length scale ξ . The action for such fluctuation is $S_\xi = \int dt [\frac{1}{2} \xi^d (\dot{\phi})^2 - \frac{1}{2} \xi^{d-2} \phi^2 - \frac{1}{4} b \xi^d \phi^4]$
→ Oscillator with mass $M = \xi^d$ and spring constant $K = \xi^{d-2}$.
Oscillator frequency $\omega = \sqrt{K/M} = 1/\xi$.
Potential energy for quantum fluctuation $E = \frac{1}{2} \omega = \frac{1}{2} \xi^{d-2} \phi^2$.
Fluctuation $\phi^2 = \xi^{1-d}$.
- Compare $\xi^{d-2} \phi^2$ and $b \xi^d \phi^4$: $\frac{b \xi^d \phi^4}{\xi^{d-2} \phi^2} = b \xi^{3-d}$ for $\xi \rightarrow \infty$, we conclude **the $b\phi^4$ term is irrelevant for $d > 3$. Relevant for $d < 3$**

Simple rules to test relevant/irrelevant perturbations

- After rescaling t to make $\rho = g$ and rescaling ϕ to make $\rho = g = 1$, the action becomes $S = \int dt d^d x \left[\frac{1}{2}(\dot{\phi})^2 - \frac{1}{2}(\partial_x \phi)^2 - \frac{1}{2}a\phi^2 - \frac{1}{4}b\phi^4 \right]$

- **Estimate from dimensional analysis:**

$$[S] = [L]^0 \text{ (from } e^{-iS}\text{)}. [t] = [L] \text{ (from } \frac{1}{2}(\dot{\phi})^2 - \frac{1}{2}(\partial_x \phi)^2\text{)}$$

$$[\phi] = [L]^{\frac{1-d}{2}}, [a] = L^{-2}, [b] = [L]^{d-3}$$

- **Counting dimensions:**

$$[t] = -1, [S] = 0.$$

$$[\phi] = \frac{d-1}{2}, [a] = 2, [b] = 3 - d.$$

- From the scaling dimensions, we can see that the quantum fluctuations of ϕ^2 are given by $\phi^2 \sim L^{1-d}$, and the dimensionless ratio of $L^d \frac{1}{L^2} \phi^2$ and $L^d b \phi^4$ terms is given by $\frac{bL^d \phi^4}{L^{d-2} \phi^2} \sim bL^{3-d}$

The $b\phi^4$ term is irrelevant if $[b] < 0$. Relevant if $[b] > 0$.

The $a\phi^2$ term is always relevant since $[a] = 2 > 0$.

- **More precise definition of scaling dimension:**

The correlation of ϕ at the critical point $a = b = 0$

$$\langle \phi(x)\phi(y) \rangle = \frac{1}{|x-y|^{2h_\phi}}. h_\phi \text{ is the scaling dimension of } \phi: h_\phi = \frac{d-1}{2}.$$

Specific heat at the critical point

- Thermal energy density

$$\epsilon_T = \int_{-\infty}^{+\infty} \frac{dk}{2\pi} \frac{v|k|}{e^{v|k|/k_B T} - 1} = 2 \frac{k_B^2 T^2}{2\pi v} \int_0^{+\infty} dx \frac{x}{e^x - 1} = \frac{k_B^2 T^2}{v} \frac{\pi}{6}$$

where $\int_0^{+\infty} dx \frac{x}{e^x - 1} = \frac{\pi^2}{6}$

- Specific heat

$$c_T = \frac{\partial \epsilon_T}{\partial T} = k_B \frac{k_B T}{v} \frac{\pi}{3} = \left(\frac{\pi}{6} k_B \frac{k_B T}{v} \right)_R + \left(\frac{\pi}{6} k_B \frac{k_B T}{v} \right)_L$$

- *The above result is incorrect. The correct one is*


$$c_T = \left(\frac{1}{2} \frac{\pi}{6} k_B \frac{k_B T}{v} \right)_R + \left(\frac{1}{2} \frac{\pi}{6} k_B \frac{k_B T}{v} \right)_L$$

- $\frac{1}{2} = c$ is called the **central charge** = number of modes.
- Many-body spectrum for one right-moving mode ($c = 1$):
 $1, 1, 2, 3, 5, 7, 11, \dots$ = partition number

Specific heat away from the critical point

Away from the critical point, the boson dispersion becomes

$\epsilon_k = \sqrt{v^2 k^2 + \Delta^2}$ where Δ is the many-body spectrum gap on a **ring** (the energy to create a single boson).



Many-body spectrum for a ring

many-body spectrum = spectrum of the set of the oscillators

($\times 2$ in the symmetry breaking phases)

Specific heat

$$c \sim T^\alpha e^{-\frac{\Delta}{k_B T}}$$

The above result is correct in the symmetric phase, but incorrect in the symmetry breaking phase. The correct one is

$$c \sim T^\alpha e^{-\frac{\Delta/2}{k_B T}}$$


Remark: The gap in many-body spectrum for an open line is $\Delta/2$.

What really is a quasiparticle? \rightarrow factor 1/2


The answer is very different for gapped system and gapless systems. Here, we only consider the definition of quasiparticle for gapped systems.

Consider a many-body system $H_0 = \sum_x H_x$, with ground state $|\Psi_{\text{grnd}}\rangle$.

- a point-like excitation above the ground state is a many-body wave function $|\Psi_\xi\rangle$ that has an energy bump at location ξ :
energy density = $\langle \Psi_\xi | H_x | \Psi_\xi \rangle$



More precisely, point-like excitations at locations ξ_i are something that can be trapped by local traps δH_{ξ_i} : $|\Psi_{\xi_i}\rangle$ is the gapped ground state of $H_0 + \sum_i \delta H_{\xi_i}$ – the Hamiltonian with traps.



Local and topological excitations

Consider a many-body state $|\Psi_{\xi_1, \xi_2, \dots}\rangle$ with several point-like excitations at locations ξ_j .

Can the first point-like excitation at ξ_1 be created by a local operator O_{ξ_1} from the ground state: $|\Psi_{\xi_1, \xi_2, \dots}\rangle = O_{\xi_1} |\Psi_{\xi_2, \dots}\rangle$?

$|\Psi_{\xi_1, \xi_2, \dots}\rangle$ = the ground state of $H_0 + \delta H_{\xi_1} + \delta H_{\xi_2} + \dots$

$|\Psi_{\xi_2, \dots}\rangle$ = the ground state of $H_0 + \delta H_{\xi_2} + \dots$

If yes: the point-like excitation at ξ_1 is a **local** excitation

If no: the point-like excitation at ξ_1 is a **topological** excitation


Experimental consequence of topological excitations

- The topological excitations are **fractionalized** local excitations: a spin-flip can be viewed as a bound state of two wall excitations $\text{spin-flip} = \text{wall} \otimes \text{wall}$.



- Energy cost of spin-flip $\Delta_{\text{flip}} = 4J$
Energy cost of domain wall $\Delta_{\text{wall}} = 2J$.

- The many-body spectrum gap on a ring $\Delta = \Delta_{\text{flip}} = 2\Delta_{\text{wall}}$. This gap can be measured by neutron scattering.



- The thermal activation gap measured by specific heat $c \sim T^\alpha e^{-\frac{\Delta_{\text{therm}}}{k_B T}}$ is $\Delta_{\text{therm}} = \Delta_{\text{wall}}$.

The difference of the neutron gap Δ and the thermal activation gap $\Delta_{\text{therm}} \rightarrow$ fractionalization.

Another example: 1D spin-dimer state

Consider a $SO(3)$ spin rotation symmetric Hamiltonian H_0 whose ground states are spin-dimer state formed by spin-singlets, which break the translation symmetry but not spin rotation symmetry:

$$\begin{aligned} &(\uparrow\downarrow)(\uparrow\downarrow)(\uparrow\downarrow)(\uparrow\downarrow)(\uparrow\downarrow)(\uparrow\downarrow)(\uparrow\downarrow)(\uparrow\downarrow) \\ &\downarrow(\uparrow\downarrow)(\uparrow\downarrow)(\uparrow\downarrow)(\uparrow\downarrow)(\uparrow\downarrow)(\uparrow\downarrow)(\uparrow\downarrow)(\uparrow\downarrow)(\uparrow\downarrow) \end{aligned}$$

- Local excitation = spin-1 excitation

$$(\uparrow\downarrow)(\uparrow\downarrow)(\uparrow\downarrow)\uparrow\uparrow(\uparrow\downarrow)(\uparrow\downarrow)(\uparrow\downarrow)(\uparrow\downarrow)$$

- Topo. excitation (domain wall) = spin-1/2 excitation (spinon)

$$(\uparrow\downarrow)(\uparrow\downarrow)\uparrow(\uparrow\downarrow)(\uparrow\downarrow)(\uparrow\downarrow)\uparrow(\uparrow\downarrow)(\uparrow\downarrow)$$

- Neutron scattering only creates the spin-1 excitation = two spinons. It measures the two-spinon gap (spin-1 gap).
Thermal activation sees single spinon gap.

Neutron scattering spectrum

Neutron dump energy-momentum into the sample creating a few excitations.


- Without fractionalization, nor trans. symm. breaking

$$\epsilon_{\text{spin-1}}(k) = 2.6 + 2 \cos(k)$$


- With fractionalization and trans. sym. breaking

$$\epsilon_{\text{spin-1/2}}(k) = \frac{1}{2} \epsilon(2k)_{\text{spin-1}}$$


one spin-1 + two spin-1/2



two spin-1/2 + four spin-1/2



k



k

2D Spin liquid without symmetry breaking (topo. order)

The spin-1 fractionalization into spin-1/2 spinon can happen in 2D spin liquid without translation and $SO(3)$ spin-rotation symmetry breaking:

- On square lattice:




chiral spin liquid $\sum \Psi(RVB)|RVB\rangle \rightarrow$ topological order

Kalmeyer-Laughlin PRL **59** 2095 (87); Wen-Wilczek-Zee PRB **39** 11413 (89)

Z_2 spin liquid $\sum |RVB\rangle$ (emergent low energy Z_2 gauge theory)

Read-Sachdev PRL **66** 1773 (91); Wen PRB **44** 2664 (91)


Z_2 -charge (spin-1/2) = Spinon. Z_2 -vortex (spin-0) = Vison. Bound state = fermion (spin-1/2).




2D Spin liquid without symmetry breaking (topo. order)

- On Kagome lattice:

Feng et al arXiv:1702.01658 $\text{Cu}_3\text{Zn}(\text{OH})_6\text{FBr}$



© American Physical Society. All rights reserved. This content is excluded from our Creative Commons license. For more information, see <https://ocw.mit.edu/help/faq-fair-use>.




© IOP Publishing. All rights reserved. This content is excluded from our Creative Commons license. For more information, see <https://ocw.mit.edu/help/faq-fair-use>.

J_1 - J_2 - J_3 model Gong-Zhu-Balents-Sheng arXiv:1412.1571

- Uniform spin susceptibility comes from spin excitations:

$\chi \sim e^{-\Delta_{\text{spinon}}/k_B T}$. In a strong magnetic field, the activation gap Δ_{spinon} is reduced to $\Delta_{\text{spinon}} - Bgs$.

Knowing the g -factor, we can measure the spin s of the spinons.



Duality between 1D boson/spin and 1D fermion systems

To obtain the correct critical theory for the transverse Ising model, we need to use the duality between 1D boson/spin systems and 1D fermion systems.

Duality: Two different theories that describe the same thing.
Two different looking theories that are equivalent.

- If we view down-spin as vacuum and up-spin as a boson, we can view a hard-core boson system as a spin-1/2 system. Now we view a system of hard-core bosons hopping on a line/ring of L sites as a spin-1/2 system. How to write down the spin Hamiltonian to describe such a boson-hopping system?

$\sigma_i^\pm = (\sigma_i^x \pm i\sigma_i^y)/2$: σ_i^- annihilates (σ_i^+ creates) a boson at site- i , $|\downarrow\rangle = |0\rangle, |\uparrow\rangle = |1\rangle$. $H_{\text{boson-hc}} = \sum_i (-t\sigma_i^+ \sigma_{i+1}^- + h.c.)$ describes a hard-core bosons hopping model.

- Similarly, we can also view a system of spin-less fermions on a line/ring of L sites as a spin-1/2 system. How to write down the spin Hamiltonian for such a fermion-hopping system?

Jordan-Wigner transformation on a 1D line of L sites

- $c_i = \sigma_i^+ \prod_{j<i} \sigma_j^z$, $\sigma^\pm = (\sigma^x \pm i\sigma^y)/2$. One can check that
$$\{c_i, c_j\} = \{c_i^\dagger, c_j^\dagger\} = 0, \quad \{c_i, c_j^\dagger\} = \delta_{ij}, \quad \{A, B\} \equiv AB - BA.$$
$$c_i^\dagger, c_i \text{ create/annihilate a fermion at site-}i, \quad |\downarrow\rangle = |0\rangle, \quad |\uparrow\rangle = |1\rangle$$
- Mapping between spin/boson chain and fermion chain:
$$c_i^\dagger c_i = \sigma_i^- \sigma_i^+ = (-\sigma_i^z + 1)/2 = n_i, \quad \text{fermion number operator}$$
$$c_i^\dagger c_{i+1} = \sigma_i^- \sigma_{i+1}^+ \sigma_i^z = \sigma_i^- \sigma_{i+1}^+, \quad c_i c_{i+1} = \sigma_i^+ \sigma_{i+1}^+ \sigma_i^z = -\sigma_i^+ \sigma_{i+1}^+$$
- **XY model = fermion model** on an open chain
$$H_{\text{fermion}} = \sum_i (-t c_i^\dagger c_{i+1} + h.c.) - \mu n_i \quad \leftrightarrow$$
$$H_{\text{XY}} = \sum_i (-t \sigma_i^+ \sigma_{i+1}^- + h.c.) + \mu \frac{\sigma_i^z}{2} = \sum_i -\frac{t}{2} (\sigma_i^x \sigma_{i+1}^x + \sigma_i^y \sigma_{i+1}^y) + \mu \frac{\sigma_i^z}{2}$$
- A phase transition in XY model: as we tune μ through $\mu_c = \pm 2t$, the ground state energy density ϵ_μ has a singularity
→ a phase transition.

How to solve the model exactly to obtain the above result?

The model H_{fermion} or H_{XY} looks not solvable since H 's are not a sum of commuting terms.

Make the Hamiltonian into a sum of commuting terms

- The anti-commutation relation

$$\{c_i, c_j\} = \{c_i^\dagger, c_j^\dagger\} = 0, \quad \{c_i, c_j^\dagger\} = \delta_{ij}$$

is invariant under the unitary transformation of the fermion operators:

$$\tilde{c}_i = U_{ij} c_j : \quad \{\tilde{c}_i, \tilde{c}_j\} = \{\tilde{c}_i^\dagger, \tilde{c}_j^\dagger\} = 0, \quad \{\tilde{c}_i, \tilde{c}_j^\dagger\} = \delta_{ij}$$

- Assume the fermions live on a ring. see the homework

Let $\psi_k = \frac{1}{\sqrt{L}} \sum_i e^{ikj} c_j$ ($k = \frac{2\pi}{L} \times \text{integer}$)

$$H_{\text{fermion}} = \sum_i (-tc_i^\dagger c_{i+1} + h.c.) + gc_i^\dagger c_i = \sum_k \epsilon(k) \psi_k^\dagger \psi_k$$

$$\epsilon(k) = -2t \cos k - \mu, \quad [\psi_k^\dagger \psi_k, \psi_{k'}^\dagger \psi_{k'}] = 0, \quad n_k \equiv \psi_k^\dagger \psi_k = \pm 1.$$

- From the one-body dispersion, we obtain many-body energy spectrum

$$E = \sum_k \epsilon(k) n_k, \quad K = \sum_k kn_k \bmod \frac{2\pi}{a}, \quad n_k = 0, 1.$$

Majorana fermions and critical point of Ising model

- $\lambda_i^x = \sigma_i^x \prod_{j<i} \sigma_j^z$, $\lambda_i^y = \sigma_i^y \prod_{j<i} \sigma_j^z$. One can check that

$$(\lambda_i^x)^\dagger = \lambda_i^x, (\lambda_i^y)^\dagger = \lambda_i^y; \quad \{\lambda_i^x, \lambda_j^x\} = \{\lambda_i^y, \lambda_j^y\} = 2\delta_{ij}, \quad \{\lambda_i^x, \lambda_j^y\} = 0.$$

- **Ising model = Majorana-fermion** on a open chain of L sites:


$$\lambda_i^x \lambda_j^y = i\sigma_i^z, \quad \lambda_i^y \lambda_{i+1}^x = \sigma_i^y \sigma_{i+1}^x \sigma_i^z = i\sigma_i^x \sigma_{i+1}^x$$

$$H_{\text{Ising}} = \sum_i -\sigma_i^x \sigma_{i+1}^x - h\sigma_i^z \leftrightarrow H_{\text{fermion}} = \sum_i i\lambda_i^y \lambda_{i+1}^x + ih\lambda_i^x \lambda_i^y$$

Critical point (gapless point) is at $h = 1$ (not $h = 2$ from meanfield theory): $H_{\text{fermion}}^{\text{critical}} = \sum_i i\eta_l \eta_{l+1}$, $\eta_{2i+1} = \lambda_i^x$, $\eta_{2i} = \lambda_i^y$.

- In k -space, $\psi_k = \frac{1}{\sqrt{2}} \sum_l \frac{e^{i\frac{k}{2}l}}{\sqrt{2L}} \eta_l$, $\frac{k}{2} = \frac{2\pi}{2L} n \in [-\pi, \pi]$:


$$\psi_k^\dagger = \psi_{-k}, \quad \{\psi_k^\dagger, \psi_{k'}\} = \delta_{k-k'} \quad (\text{assume on a ring})$$



$$H_{\text{fermion}}^{\text{critical}} = \sum_{k \in [-2\pi, 2\pi]} 2i e^{i\frac{1}{2}k} \psi_{-k} \psi_k = \sum_{k \in [0, 2\pi]} \epsilon(k) \psi_k^\dagger \psi_k, \quad \epsilon(k) = 4 \left| \sin \frac{k}{2} \right|.$$

1D Ising critical point: 1/2 mode of right (left) movers

- The Majorana fermion contain a right-moving mode $\epsilon = vk$ and a left-moving modes. $\epsilon = -vk$



- Thermal energy density (for a right moving mode):

$$\epsilon_T = \int_0^{+\infty} \frac{dk}{2\pi} \frac{vk}{e^{vk/k_B T} + 1} = \frac{k_B^2 T^2}{2\pi v} \int_0^{+\infty} dx \frac{x}{e^x + 1} = \frac{k_B^2 T^2}{v} \frac{\pi}{24}$$

where $\int_0^{+\infty} dx \frac{x}{e^x + 1} = \frac{\pi^2}{12}$

- Specific heat

$$c_T = \frac{\partial \epsilon_T}{\partial T} = \frac{1}{2} k_B \frac{k_B T}{v} \frac{\pi}{6}$$

Central charge $c = 1/2$ for right (left) movers.

- On a ring of size L and at critical point: the ground state energy has a form $E = \epsilon L + \frac{2\pi v}{L} (-\frac{c}{24})$, where c in the “Casimir term” (the $1/L$ term) is also the central charge.

Do we have a similar result for an open Line?

A story about central charge c (conformal field theory)

- Central charge is a property of 1D gapless system with a finite and unique velocity. $c = c_L + c_R = 0$ for gapped systems.
- It has an additive property: $A \boxtimes_{\text{stacking}} B = C \rightarrow c_A + c_B = c_C$
- It measures how many low energy excitations are there.
Specific heat (heat capacity per unit length) $C = c \frac{\pi}{6} \frac{T}{v}$

A story about central charge c (conformal field theory)

- Central charge is a property of 1D gapless system with a finite and unique velocity. $c = c_L + c_R = 0$ for gapped systems.
- It has an additive property: $A \boxtimes_{\text{stacking}} B = C \rightarrow c_A + c_B = c_C$
- It measures how many low energy excitation are there.
Specific heat (heat capacity per unit length) $C = c \frac{\pi}{6} \frac{T}{v}$
- Why ground state energy $E = \rho_\epsilon L - \frac{c}{24} \frac{2\pi}{L}$ sees central charge ($v = 1$)?
Partition function: $Z(\beta, L) = \text{Tr}(e^{-\beta H}) = e^{-\beta L \rho_\epsilon - \frac{2\pi\beta}{L} \frac{c}{24}} \Big|_{\beta \rightarrow \infty}$
- A magic: emergence of $O(2)$ symmetry in space-(imaginary-)time

$$Z(\beta, L) = Z(L, \beta), \quad \text{have used } v = 1.$$

This allows us to find $Z(\beta, L) = e^{-\beta L \rho_\epsilon - \frac{2\pi L}{\beta} \frac{c}{24}} \Big|_{L \rightarrow \infty}$

$$\begin{aligned} \text{Free energy density } f &= \rho_\epsilon - \frac{2\pi}{(\beta)^2} \frac{c}{24} \\ &= \rho_\epsilon - 2\pi T^2 \frac{c}{24} \end{aligned}$$

$$\text{Specific heat } C = -T \frac{\partial^2 F}{\partial T^2} = T \frac{\pi}{6} c$$

The neutron scattering and spectral function (Ising model)

Assume the neutron spin couples to Ising spin via $S_i^z \sim \sigma_i^z$ (no S^z -spin flip, but scattering flips $S^{x,y}$). After scattering, the neutron dump something to the system $|\Psi\rangle \rightarrow \sigma_i^z |\Psi\rangle$. What is the scattering spectrum? **The spectra function of σ_i^z :**

$$I(E, K) = \langle \Psi | \sigma_i^z \delta(\hat{H} - E) \delta(\hat{K} - K) \sigma_i^z | \Psi \rangle$$

$$\sigma_i^z = i\eta_{2i}\eta_{2i+1} = \frac{2i}{L} \sum_{k_1, k_2} e^{ik_1 i} e^{ik_2(i+\frac{1}{2})} \psi_{k_1} \psi_{k_2}$$

$$\begin{aligned} I(E, K) &= \frac{4}{L^2} \langle \Psi | \sum_{k_1, k_2} e^{ik_1 i} e^{ik_2(i+\frac{1}{2})} \psi_{k_1} \psi_{k_2} \delta(\epsilon_{k'_1} + \epsilon_{k'_2} - E) \\ &\quad \delta(k'_1 + k'_2 - K) \sum_{k'_1, k'_2} e^{-ik'_1 i} e^{-ik'_2(i+\frac{1}{2})} \psi_{k'_2}^\dagger \psi_{k'_1}^\dagger | \Psi \rangle \\ &= \frac{4}{L^2} \sum_{k_1, k_2 \in [0, 2\pi]} \delta(\epsilon_{k_1} + \epsilon_{k_2} - E) \delta(k_1 + k_2 - K) (1 - e^{i\frac{1}{2}(k_1 - k_2)}) \end{aligned}$$


The neutron scattering and spectral function (Ising model)

$$I(E, K) = 4 \int_0^{2\pi} \frac{dk_1 dk_2}{(2\pi)^2} \delta(\epsilon_{k_1} + \epsilon_{k_2} - E) \delta(k_1 + k_2 - K) (1 - \cos \frac{k_1 - k_2}{2})$$


$$I_0(E, K) = 4 \int_0^{2\pi} \frac{dk_1 dk_2}{(2\pi)^2} \delta(\epsilon_{k_1} + \epsilon_{k_2} - E) \delta(k_1 + k_2 - K)$$

where $\epsilon_k = 4|\sin \frac{k}{2}|$.

$I(E, K)$



$I_0(E, K)$: two-fermion density of states



$-\pi$ K π

$-\pi$ K π

- What is the spectral function for σ_i^x ? for $\sigma_i^x \sigma_j^x$? Why σ_i^x is hard?



A general picture of spectre function

We can understand the spectral function of an operator O_x by writing it in terms of quasiparticle creating/annihilation operators

$$\begin{aligned} O_i &= C_1 a_i^\dagger + C_2 a_i^\dagger a_{i+1}^\dagger + \dots \\ &= C_1 \int dk a_k^\dagger + C_2 \int dk_1 dk_2 a_{k_1}^\dagger a_{k_2}^\dagger e^{-i[k_1 i + k_2 (i+1)]} + \dots \end{aligned}$$

Assume one-particle spectrum to be $\epsilon(k) = 2.6 + 2 \cos(k) \rightarrow$

Two-particle spectrum will be $E = \epsilon(k_1) + \epsilon(k_2)$, $K = k_1 + k_2$



Spectre function and time-ordered correlation functions

- Consider a 0d system with ground state $|0\rangle$ with energy $E_0 = 0$. An operator O creates excitations, and have a spectral function

$$I(\omega) = \langle 0|O^\dagger \delta(\hat{H} - \omega) O|0\rangle.$$

- Time-ordered correlation function of $O(t) = e^{i\hat{H}t} O e^{-i\hat{H}t}$:

$$\begin{aligned} G(t) &= i \langle 0|\mathcal{T}[O(t)O(0)]|0\rangle = i \begin{cases} \langle 0|O(t)O(0)|0\rangle, & t > 0 \\ \langle 0|O(0)O(t)|0\rangle, & t < 0 \end{cases} \\ &= i \begin{cases} \langle 0|O e^{-i\hat{H}t} O|0\rangle, & t > 0 \\ \langle 0|O e^{i\hat{H}t} O|0\rangle, & t < 0 \end{cases} = i \begin{cases} \int_0^{+\infty} d\omega e^{-i\omega t} I(\omega), & t > 0 \\ \int_0^{+\infty} d\omega e^{i\omega t} I(\omega), & t < 0 \end{cases} \end{aligned}$$

$$\begin{aligned} G(\omega) &= \int dt G(t) e^{i\omega t} = i \int_0^{+\infty} dt \int_0^{+\infty} d\omega' \left(e^{-i(\omega' - \omega - i0^+)t} I(\omega') - e^{-i(\omega' + \omega - i0^+)t} I(\omega') \right) \\ &= \int_0^{+\infty} d\omega' \left(\frac{I(\omega')}{\omega' - \omega - i0^+} - \frac{I(\omega')}{\omega' + \omega - i0^+} \right) = \int_{-\infty}^{+\infty} d\omega' \frac{I(|\omega'|)}{\omega' - \omega - i0^+ \operatorname{sgn}\omega'} \end{aligned}$$

$$I(\omega) = \frac{\operatorname{sgn}(\omega)}{\pi} \operatorname{Im} G(\omega). \quad \text{Adding } i0^+ \text{ to regulate the integral } \int_0^{+\infty} dt$$

- In higher dimensions: $G(t, \mathbf{x}) \rightarrow G(\omega, \mathbf{k}) \rightarrow I(\omega, \mathbf{k}) = \frac{\operatorname{sgn}(\omega)}{\pi} \operatorname{Im} G(\omega, \mathbf{k})$

The neutron scattering and spectral function (XY model)

1D XY model (superfluid of bosons) = 1D non-interacting fermions

$$H_{XY} = \sum_i -\frac{t}{2}(\sigma_i^x \sigma_{i+1}^x + \sigma_i^y \sigma_{i+1}^y) - \mu \frac{\sigma_i^z}{2} \leftrightarrow H_f = \sum_i (t c_i^\dagger c_{i+1} + h.c.) - \mu n_i$$

Let us assume the neutron coupling is $S_i^z \sim \sigma_i^z$ (ie neutrons see the boson density) \rightarrow Spectral function of operator $\sigma_i^z = c_i^\dagger c_i$ (adding a particle-hole pair)

$$\begin{aligned} I(E, K) &= \langle \Psi | c_i^\dagger c_i \delta(\hat{H} - E) \delta(\hat{K} - K) c_i^\dagger c_i | \Psi \rangle \\ &= \frac{1}{L^2} \langle \Psi | \sum_{k_1, k_2} e^{ik_1 i} e^{ik_2 i} \psi_{k_1}^\dagger \psi_{k_2} \delta(-\epsilon_{k_1'} + \epsilon_{k_2'} - E) \\ &\quad \delta(-k_1' + k_2' - K) \sum_{k_1', k_2'} e^{-ik_1' i} e^{-ik_2' i} \psi_{k_1'}^\dagger \psi_{k_2'} | \Psi \rangle \\ &= \int_{\epsilon_{k_1} < 0, \epsilon_{k_2} > 0} \frac{dk_1 dk_2}{(2\pi)^2} \delta(-\epsilon_{k_1} + \epsilon_{k_2} - E) \delta(-k_1 + k_2 - K) \end{aligned}$$

where $\epsilon_k = 2t \cos k - \mu$ and $c_i = \frac{1}{\sqrt{L}} \sum_k e^{ik_i} \psi_k$


The neutron scattering and spectral function (XY model)

Spectral function of $n_i \sim \sigma_i^z$ for the superfluid/XY-model

For $\mu = 0$, $\langle \sigma_i^z \rangle = 0$



For $\mu = -1$, $\langle \sigma_i^z \rangle \neq 0$





Particle-hole spectral function. In addition to the low energy excitations near $k = 0$, why are there low energy excitations at large $K_{\pm} = \pm 2\pi n$? K_{\pm} only depend on boson density n ! What is the single particle spectral function of σ_i^+ ? $\sigma_i^+ = c_i^{\dagger} \prod_{j < i} (2c_j^{\dagger} c_j - 1)$

The neutron scattering and spectral function (XY model)

Particle-particle spectral function of $\sigma_i^+ \sigma_{i+1}^+$ (adding two bosons)

$$\begin{aligned} I(E, K) &= \langle \Psi | c_{i+1} c_i \delta(\hat{H} - E) \delta(\hat{K} - K) c_i^\dagger c_{i+1}^\dagger | \Psi \rangle \\ &= \frac{1}{L^2} \langle \Psi | \sum_{k_1, k_2} e^{ik_1(i+1)} e^{ik_2 i} \psi_{k_1} \psi_{k_2} \delta(\epsilon_{k_1} + \epsilon_{k_2} - E) \\ &\quad \delta(k_1' + k_2' - K) \sum_{k_1', k_2'} e^{-ik_1'(i+1)} e^{-ik_2' i} \psi_{k_2'}^\dagger \psi_{k_1'}^\dagger | \Psi \rangle \\ &= \int_{\substack{\epsilon_{k_1} > 0 \\ \epsilon_{k_2} > 0}} \frac{dk_1 dk_2}{(2\pi)^2} \delta(\epsilon_{k_1} + \epsilon_{k_2} - E) \delta(k_1 + k_2 - K) [1 - \cos(k_1 - k_2)] \end{aligned}$$

$\mu = 0$ and
 $\mu = -1$
2-particle
spectral function



XY model for superfluid: dynamical variational approach

Compute single-particle spectral function using an approximation

We are going to use the approximated variational approach for XY model (not bad for superfluid phase. See also prob. 4.2):

$$H = - \sum_i J(\sigma_i^x \sigma_{i+1}^x + \sigma_i^y \sigma_{i+1}^y) + h \sigma_i^z.$$

Trial wave function $|\Psi_{\phi_i}\rangle = \otimes_i |\phi_i\rangle$,

$$\text{where } |\phi_i\rangle = \frac{|\uparrow\rangle + \phi_i |\downarrow\rangle}{\sqrt{1 + |\phi_i|^2}}, \quad \langle \sigma_i^+ \rangle = \frac{\phi_i}{1 + |\phi_i|^2}.$$

- Average energy $\bar{H} = - \sum_i \left[2J \frac{\phi_i \phi_{i+1}^* + h.c.}{(1 + |\phi_i|^2)(1 + |\phi_{i+1}|^2)} + h \frac{1 - |\phi_i|^2}{1 + |\phi_i|^2} \right]$

Geometric phase term $\langle \phi_i | \frac{d}{dt} | \phi_i \rangle = \frac{\dot{\phi}_i^* \phi_i}{1 + |\phi_i|^2} + \frac{d}{dt} \#$

Phase space Lagrangian in symmetry breaking phase (up to φ_i^2)

($\phi_i = \bar{\phi} + \varphi_i$ for $J > 0$ or $\phi_i = \bar{\phi}(-)^i + \varphi_i$ for $J < 0$)

$$L = \langle \Phi_{\phi_i} | i \frac{d}{dt} - H | \Phi_{\phi_i} \rangle = \sum_i i \phi_i^* \dot{\phi}_i + 2J(\phi_i \phi_{i+1}^* + h.c.) - 2h|\phi_i|^2 - g|\phi_i|^4$$

$$= \sum_i i \varphi_i^* \dot{\varphi}_i + 2J(\varphi_i \varphi_{i+1}^* + h.c.) - 2h\varphi_i \varphi_i^* - g\bar{\phi}^2 [4\varphi_i \varphi_i^* + \underbrace{\varphi_i^2 + (\varphi_i^*)^2}_{\text{symm breaking}}]$$

with $g\bar{\phi}^2 = 2|J| - h$.

Quantum XY model

Quantization:

$$[\varphi_i, \varphi_j^\dagger] = \delta_{ij}, \quad \varphi_i = \frac{1}{\sqrt{L}} \sum_k e^{iki} a_k, \quad [a_k, a_q^\dagger] = \delta_{kq}$$

$$H = \sum_i -2J(\varphi_i \varphi_{i+1}^\dagger + h.c.) + 2h \varphi_i^\dagger \varphi_i + (2|J| - h)(4\varphi_i^\dagger \varphi_i + \varphi_i \varphi_i + \varphi_i^\dagger \varphi_i^\dagger)$$

$$= \sum_k (-4J \cos k + 8|J| - 2h) a_k^\dagger a_k + (2|J| - h)(a_k a_{-k} + a_k^\dagger a_{-k}^\dagger)$$

$$= \sum_{k \in [0, \pi]} \begin{pmatrix} a_k^\dagger & a_{-k} \end{pmatrix} \begin{pmatrix} -4J \cos k + 8|J| - 2h & 2(2|J| - h) \\ 2(2|J| - h) & -4J \cos k + 8|J| - 2h \end{pmatrix} \begin{pmatrix} a_k \\ a_{-k}^\dagger \end{pmatrix}$$

$$= \sum_{k \in [0, \pi]} \begin{pmatrix} a_k^\dagger & a_{-k} \end{pmatrix} \begin{pmatrix} \epsilon_k & \Delta \\ \Delta & \epsilon_k \end{pmatrix} \begin{pmatrix} a_k \\ a_{-k}^\dagger \end{pmatrix}, \quad \begin{aligned} \epsilon_k &= -4J \cos k + 8|J| - 2h, \\ \Delta &= 2(2|J| - h). \end{aligned}$$

To diagonalize the above Hamiltonian, let

$$\begin{pmatrix} a_k \\ a_{-k}^\dagger \end{pmatrix} = U \begin{pmatrix} b_k \\ b_{-k}^\dagger \end{pmatrix}, \quad U = \begin{pmatrix} u_k & -v_k \\ -v_k & u_k \end{pmatrix}, \quad U \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} U = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

where $u_k^2 - v_k^2 = 1$

Quantum XY model

$$H = \sum_{k \in [0, \pi]} \begin{pmatrix} a_k^\dagger & a_{-k} \end{pmatrix} \begin{pmatrix} \epsilon_k & \Delta \\ \Delta & \epsilon_k \end{pmatrix} \begin{pmatrix} a_k \\ a_{-k}^\dagger \end{pmatrix}$$

$$U \begin{pmatrix} \epsilon_k & \Delta \\ \Delta & \epsilon_k \end{pmatrix} U = \begin{pmatrix} (u^2 + v^2)\epsilon_k - 2uv\Delta & (u^2 + v^2)\Delta - 2uv\epsilon_k \\ (u^2 + v^2)\Delta - 2uv\epsilon_k & (u^2 + v^2)\epsilon_k - 2uv\Delta \end{pmatrix} \\ = \begin{pmatrix} E_k & 0 \\ 0 & E_k \end{pmatrix}, \quad E_k = \sqrt{\epsilon_k^2 - \Delta^2}$$

$$u^2 + v^2 = \frac{\epsilon_k}{E_k}, \quad 2uv = \frac{\Delta}{E_k},$$

$$u = \sqrt{\frac{\frac{\epsilon_k}{E_k} + 1}{2}}, \quad v = \sqrt{\frac{\frac{\epsilon_k}{E_k} - 1}{2}}$$


$$H = \sum_k b_k^\dagger \underbrace{\sqrt{(-4J \cos k + 8|J| - 2h)^2 - (4|J| - 2h)^2}}_{\sqrt{\epsilon_k^2 - \Delta^2} = E_k \rightarrow 0|_{k \rightarrow 0}, \text{ spin-wave dispersion}} b_k$$

The spectral function – XY model (only for $\langle \sigma^+ \rangle = \bar{\phi}$)

- Spectral function for $\sigma^+ \sim \bar{\phi} + \varphi_i^\dagger$,
and $(\sigma^+)^2 \sim \bar{\phi}^2 + 2\bar{\phi}\varphi_i^\dagger + (\varphi_i^\dagger)^2$

$$\begin{aligned}\varphi_i^\dagger &= \frac{1}{\sqrt{L}} \sum_k e^{-iki} a_k^\dagger \\ &= \frac{1}{\sqrt{L}} \sum_k e^{-iki} (u_k b_k^\dagger - v_k b_{-k})\end{aligned}$$


$$I(E, K) \sim u_K^2 \delta(E_K - E) = \frac{\frac{\epsilon_k}{E_k} + 1}{2} \delta(E_K - E) \rightarrow \infty |_{k \rightarrow 0}$$



- Spectral function for $n_i = \frac{\sigma_i^z - 1}{2} \sim \sigma_i^x \sim \varphi_i + \varphi_i^\dagger$


$$\begin{aligned}\varphi_i + \varphi_i^\dagger &= \frac{1}{\sqrt{L}} \sum_k e^{-iki} (a_{-k} + a_k^\dagger) \\ &= \frac{1}{\sqrt{L}} \sum_k e^{-iki} (u_k b_{-k} - v_k b_k^\dagger + u_k b_k^\dagger - v_k b_{-k})\end{aligned}$$

$$I(E, K) \sim (u_K - v_K)^2 \delta(E_K - E) = \frac{E_k}{\epsilon_k + \Delta} \delta(E_K - E) \rightarrow 0 |_{k \rightarrow 0}$$




The spectral function – XY model (only for $\langle \sigma^+ \rangle = \bar{\phi}$)

The following picture work in higher dimension since $\langle \sigma_i^+ \rangle = \bar{\phi}$
(symmetry breaking) $\langle \sigma_i^+ \sigma_j^- \rangle \sim \text{const.}$ for large $|i - j|$



K



But does not quite work in 1 dimension (or 1+1 dimensions) since $\langle \sigma_i^+ \rangle = 0$ (no symmetry breaking).


We only have a **nearly symmetry breaking**

$$\langle \sigma_i^+ \sigma_j^- \rangle \sim \frac{1}{|i - j|^\alpha} \text{ for large } |i - j|$$

Neutron scattering spectrum for 2-dimensional α - RuCl_3

Banerjee et al arXiv:1706.07003

- Spin-1/2 on honeycomb lattice with strong spin-orbital coupling.
- Spin ordered phase below 8T, spin liquid above 8T
- Magnetic field:
 - (a-e) $B : 0, 2, 4, 6, 8\text{ T}$
 - (a-e) $T = 2\text{ K}$
 - (f) $T = 2\text{ K}, B = 0\text{ T}$



© Springer Nature Limited. All rights reserved. This content is excluded from our Creative Commons license. For more information, see <https://ocw.mit.edu/help/faq-fair-use>.

1d field theory to study no $U(1)$ symmetry breaking in 1D

Phase space Lagrangian in “symmetry breaking phase” of 1D XY model: $\phi_i = (\bar{\phi} + q_i)e^{-i\theta_i}$, $\bar{\phi}^2 = \frac{2J-h}{g}$, near the transition $\bar{\phi} \sim 0$

$$\begin{aligned} L &= \sum_i i\phi_i^* \dot{\phi}_i + 2J(\phi_i \phi_{i+1}^* + h.c.) - 2h|\phi_i|^2 - g|\phi_i|^4 \\ &\approx \sum_i (\bar{\phi} + q_i)^2 \dot{\theta}_i + \frac{1}{2} \partial_t (\bar{\phi} + q_i)^2 \\ &\quad + 2J|\bar{\phi}|^2 (e^{i(\theta_i - \theta_{i+1})} + h.c.) - 4(2J - h)q_i^2, \end{aligned}$$

where we kept up to q_i^2 terms. The total derivative term $\frac{1}{2} \partial_t (\bar{\phi} + q_i)^2$ can be dropped. The total “derivative” term $\bar{\phi}^2 \dot{\theta}_i$ cannot be dropped since it is not a total derivative $\bar{\phi}^2 \dot{\theta}_i = i\bar{\phi}^2 e^{i\theta} \partial_t e^{-i\theta}$.

1d field theory to study no $U(1)$ symmetry breaking in 1D

After dropping $q_i^2 \dot{\theta}_i$ term, we obtain

$$\begin{aligned} L &= \sum_i (\bar{\phi}^2 + 2\bar{\phi}q_i)\dot{\theta}_i - 2J|\bar{\phi}|^2(\theta_i - \theta_{i+1})^2 - 4(2J - h)q_i^2 \\ &= \int dx [\bar{\phi}^2 + \underbrace{\frac{2\bar{\phi}}{a}q(x)}_{\partial_x \varphi / 2\pi}] \dot{\theta}(x) - 2J|\bar{\phi}|^2 a [\partial_x \theta(x)]^2 - \frac{4(2J - h)}{a} q^2(x) \\ &= \int dx \frac{1}{2\pi} \partial_x \varphi \partial_t \theta - \frac{1}{4\pi} V_1 (\partial_x \theta)^2 - \frac{1}{4\pi} V_2 (\partial_x \varphi)^2 + \frac{\bar{\phi}^2}{a} \partial_t \theta \end{aligned}$$

where $V_1 = \frac{8\pi J(2J-h)a}{g}$, $V_2 = \frac{ga}{\pi}$.

- Momentum of uniform $\theta(x)$: $\int dx \frac{\partial_x \varphi}{2\pi} = \frac{\Delta \varphi}{2\pi} = \text{int.}$ $\rightarrow \varphi$ also live on S^1 :
 $\varphi \sim \varphi + 2\pi$

Both θ and φ are compact angular fields living on S^1 .

1d field theory with two angular fields

- Let $\varphi_1 = \theta$ and $\varphi_2 = \varphi$, we can rewrite that above as phase space Lagrangian as

$$L = \int dx \frac{2}{4\pi} \partial_x \varphi_2 \partial_t \varphi_1 - \frac{1}{4\pi} V_1 (\partial_x \varphi_1)^2 - \frac{1}{4\pi} V_2 (\partial_x \varphi_2)^2 + \frac{\bar{\varphi}^2}{a} \partial_t \varphi_1,$$

which has the following general form

$$L = \int dx \frac{K_{IJ}}{4\pi} \partial_x \varphi_I \partial_t \varphi_J - \frac{V_{IJ}}{4\pi} \partial_x \varphi_I \partial_x \varphi_J, \varphi_I \sim \varphi_I + 2\pi, K = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$

- A very generic 1+1D bosonic model:** Compact fields $\phi_I \sim \phi_I + 2\pi$. V is symmetric and positive definite. K is a symmetric integer matrix.
 - Positive eigenvalues of $K \rightarrow$ left movers.
 - Negative eigenvalues of $K \rightarrow$ right movers. (See next page)
 - The model is **chiral** if the right and left movers are not symmetric.
 - For bosonic system, the diagonal of K are all even. For fermionic system, some diagonal of K are odd even.
 - The field theory is not realizable by lattice model if $K \not\cong \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$, ie has gravitational anomalies.

1d field theory: right movers and left movers

- Introduce $\begin{pmatrix} \theta \\ \varphi \end{pmatrix} = U \begin{pmatrix} \phi_1 \\ \phi_2 \end{pmatrix}$, we can diagonaliz K, V simultaneously:

$$K \rightarrow U^\top K U, \quad V \rightarrow U^\top V U. \quad \text{Let } U = U_1 U_2.$$

- We first use U_1 to transform $V \rightarrow U_1^\top V U_1 = \text{id}$. $K \rightarrow U_1^\top K U_1$.

- We then use orthorgonal U_2 to transform

$$U_1^\top K U_1 \rightarrow U_2^\top U_1^\top K U_1 U_2 = \text{Diagonal}(\kappa_1, -\kappa_2, \dots) \text{ and}$$

$$U_1^\top V U_1 = \text{id} \rightarrow U_2^\top U_1^\top V U_1 U_2 = \text{id}.$$

- For our case of $K = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$, we find $U = \begin{pmatrix} (2V_1)^{-1/2} & (2V_1)^{-1/2} \\ (2V_2)^{-1/2} & -(2V_2)^{-1/2} \end{pmatrix}$.

$$K \rightarrow \begin{pmatrix} \kappa & 0 \\ 0 & -\kappa \end{pmatrix}, \quad \kappa = (V_1 V_2)^{-1/2}, \quad V \rightarrow \text{id}, \text{ and}$$

$$L = \int dx \left[\frac{1}{2\pi} \partial_x \varphi \partial_t \theta - \frac{1}{4\pi} V_1 (\partial_x \theta)^2 - \frac{1}{4\pi} V_2 (\partial_x \varphi)^2 + \underbrace{\frac{\bar{\phi}^2}{a}}_{\text{dropped}} \partial_t \theta \right]$$

$$= \int dx \left[\frac{1}{4\pi} (\kappa \partial_x \phi_1 \partial_t \phi_1 - \partial_x \phi_1 \partial_x \phi_1) + \frac{1}{4\pi} (-\kappa \partial_x \phi_2 \partial_t \phi_2 - \partial_x \phi_2 \partial_x \phi_2) \right]$$

- ϕ_1 and ϕ_2 are not really decoupled, since their compactness are mixed.

1d field theory – chiral boson model

$$L = \int dx \frac{\kappa}{4\pi} \partial_x \phi_1 (\partial_t \phi_1 - v \partial_x \phi_1) - \frac{\kappa}{4\pi} \partial_x \phi_2 (\partial_t \phi_2 + v \partial_x \phi_2)$$

EOM: $\partial_t \phi_1 - v \partial_x \phi_1 = 0$ and $\partial_t \phi_2 + v \partial_x \phi_2 = 0$ ($v = 1/\kappa$)

→ $\phi_1(x + vt)$ is left-mover. $\phi_2(x - vt)$ is right-mover.

- Consider only right-movers ($\phi(x) = \sum_n e^{-ikx} \phi_n$, $k = k_0 n$, $k_0 = \frac{2\pi}{L}$)

$$\begin{aligned} L &= - \int dx \frac{\kappa}{4\pi} \partial_x \phi (\partial_t \phi + v \partial_x \phi) \quad (\text{consider only } n \neq 0 \text{ terms}) \\ &= \sum_{n \neq 0} -\frac{\kappa L}{4\pi} (-ik) \phi_n (\dot{\phi}_{-n} + ivk \phi_{-n}) = \sum_{n > 0} in\kappa \phi_n (\dot{\phi}_{-n} + ivk \phi_{-n}) \end{aligned}$$

Quantize $[x, p] = i$: $[\phi_{-n}, in\kappa \phi_n] = i$, $H = \sum_{n > 0} vk n \kappa \phi_n \phi_{-n}$

Let $a_n^\dagger = \sqrt{n\kappa} \phi_n \rightarrow a_n = \sqrt{n\kappa} \phi_{-n}$

$$[a_n, a_n^\dagger] = 1, \quad H = \sum_{n > 0} vk \frac{a_n^\dagger a_n + a_n a_n^\dagger}{2} = \sum_{n > 0} vk (a_n^\dagger a_n + \frac{1}{2}).$$

Time-ordered correlation function

- Equal time correlation $\langle 0|O(x)O(y)|0\rangle \equiv \langle O(x)O(y)\rangle$
- Time dependent operator $O(t) = e^{iHt}Oe^{-iHt}$ so that

$$\langle \Phi'|O(t)|\Phi\rangle = \langle \Phi'(t)|O|\Phi(t)\rangle,$$

where $|\Phi(t)\rangle = e^{-iHt}|\Phi\rangle$, $|\Phi'(t)\rangle = e^{-iHt}|\Phi'\rangle$. We find

$$\begin{aligned} a_n^\dagger(t) &= e^{ivkt}a_n^\dagger, & \phi_n(t) &= e^{ivkt}\phi_n, \\ \phi(x, t) &= \sum_n e^{-ik(x-vt)}\phi_n, & k &= \frac{2\pi}{L}n. \end{aligned}$$

- Time-ordered correlation

$$-iG(x-y, t) = \langle \mathcal{T}[\phi(x, t)\phi(y, 0)] \rangle = \begin{cases} \langle \phi(x, t)\phi(y, 0) \rangle, & t > 0 \\ \langle \phi(y, 0)\phi(x, t) \rangle, & t < 0 \end{cases}$$

For anti-commuting operators (to make $G(x, t)$ a continuous function of x, t away from $(x, t) = (0, 0)$)

$$-iG(x-y, t) = \langle \mathcal{T}[\psi(x, t)\tilde{\psi}(y, 0)] \rangle = \begin{cases} \langle \psi(x, t)\tilde{\psi}(y, 0) \rangle, & t > 0 \\ -\langle \tilde{\psi}(y, 0)\psi(x, t) \rangle, & t < 0 \end{cases}$$

Time ordered correlation function of chiral field $\phi(x, t)$

- For $t > 0$ ($k = nk_0$, $k_0 = \frac{2\pi}{L}$)

$$\begin{aligned}\langle \phi(x, t)\phi(0, 0) \rangle &= \sum_{n_1, n_2} e^{-ik_1(x-vt)} \langle \phi_{n_1}\phi_{n_2} \rangle = \sum_{n_2 > 0} e^{ik_2(x-vt)} \underbrace{\langle \phi_{-n_2}\phi_{n_2} \rangle}_{\frac{a_{n_2}}{\sqrt{n_2\kappa}} \frac{a_{n_2}^\dagger}{\sqrt{n_2\kappa}}} \\ &= \sum_{n=1}^{\infty} e^{i2\pi \frac{x-vt}{L} n} \frac{1}{n\kappa} = -\frac{1}{\kappa} \log(1 - e^{i2\pi \frac{x-vt}{L}})\end{aligned}$$

since $\sum_{n=1}^{\infty} e^{\alpha n} \frac{1}{n} = -\log(1 - e^\alpha)$, $\text{Re}(\alpha) < 0$.

- For $t < 0$

$$\begin{aligned}\langle \phi(0, 0)\phi(x, t) \rangle &= \sum_{n_1, n_2} e^{-ik_1(x-vt)} \langle \phi_{n_2}\phi_{n_1} \rangle = \sum_{n_1 > 0} e^{-ik_1(x-vt)} \langle \phi_{-n_1}\phi_{n_1} \rangle \\ &= \sum_{n=1}^{\infty} e^{-i2\pi \frac{x-vt}{L} n} \frac{1}{n\kappa} = -\frac{1}{\kappa} \log(1 - e^{-i2\pi \frac{x-vt}{L}})\end{aligned}$$

Correlation function of vertex operator $e^{i\alpha\phi}$

- Normal ordering ($e^A e^B = e^{\frac{1}{2}[A,B]} e^{A+B}$) $[\phi_n, \phi_{-n}] = \frac{1}{\kappa n}, n > 0$

$$\begin{aligned}
 : e^{i\alpha\phi(x,t)} : &= \underbrace{e^{i\alpha \sum_{n>0} e^{ik(x-vt)} \phi_n}}_{\text{creation}} \underbrace{e^{i\alpha \sum_{n<0} e^{ik(x-vt)} \phi_n}}_{\text{annihilation}} \\
 &= e^{-\frac{\alpha^2}{2} [\sum_{n>0} e^{ik(x-vt)} \phi_n, \sum_{n<0} e^{ik(x-vt)} \phi_n]} e^{i\phi(x,t)} = \underbrace{e^{\frac{\alpha^2}{2\kappa} \sum_n \frac{1}{n}}}_{\sim (\frac{1}{a})^{\frac{\alpha^2}{2\kappa}}} e^{i\phi(x,t)}
 \end{aligned}$$

- Correlation function ($e^A e^B = e^{[A,B]} e^B e^A$)

$$\begin{aligned}
 \langle : e^{i\alpha\phi(x,t)} : : e^{-i\alpha\phi(0,0)} : \rangle &= \langle e^{i\alpha\phi_{>}(x,t)} e^{i\alpha\phi_{<}(x,t)} e^{-i\alpha\phi_{>}(0,0)} e^{-i\alpha\phi_{<}(0,0)} \rangle \\
 &= \langle e^{i\alpha\phi_{<}(x,t)} e^{-i\alpha\phi_{>}(0,0)} \rangle = \underbrace{e^{[\alpha\phi_{<}(x,t), \alpha\phi_{>}(0,0)]}}_{=e^{2\langle\phi(x,t)\phi(0,0)\rangle}} \underbrace{\langle e^{-i\alpha\phi_{>}(0,0)} e^{i\alpha\phi_{<}(x,t)} \rangle}_{=1} \\
 &= \begin{cases} (1 - e^{i2\pi \frac{x-vt+i0^+}{L}})^{-\alpha^2/\kappa}, & t > 0 \\ (1 - e^{-i2\pi \frac{x-vt-i0^+}{L}})^{-\alpha^2/\kappa}, & t < 0 \end{cases} \\
 &\approx \frac{(L/2\pi)^{\alpha^2/\kappa}}{[-i(x-vt)\text{sgn}(t) + 0^+]^{\alpha^2/\kappa}} = \frac{(L/2\pi)^{1/\kappa} e^{i\frac{1}{\kappa} \frac{\pi}{2} \text{sgn}((x-vt)t)}}{|x-vt|^{\alpha^2/\kappa}}
 \end{aligned}$$

The value of the multivalued function is in the branch of $0^+ \rightarrow +\infty$.

Correlation function of $e^{i\theta}$ and no symmetry breaking

$$\begin{aligned}
 & \langle \mathcal{T} [: e^{i\theta(x,t)} :: e^{-i\theta(0,0)} :] \rangle \quad e^{i\theta} = e^{i(\alpha\phi_1 + \alpha\phi_2)}, \alpha = (2V_1)^{-1/2} \\
 & = \langle \mathcal{T} [: e^{\frac{\alpha}{2}i\phi_1(x,t)} :: e^{-\frac{\alpha}{2}i\phi_1(0,0)} :] \rangle \langle \mathcal{T} [: e^{\frac{\alpha}{2}i\phi_2(x,t)} :: e^{-\frac{\alpha}{2}i\phi_2(0,0)} :] \rangle \\
 & = \begin{cases} (1 - e^{i2\pi \frac{-x-vt+i0^+}{L}})^{-\alpha^2/4\kappa} (1 - e^{i2\pi \frac{x-vt+i0^+}{L}})^{-\alpha^2/4\kappa}, & t > 0 \\ (1 - e^{-i2\pi \frac{-x-vt-i0^+}{L}})^{-\alpha/4\kappa} (1 - e^{-i2\pi \frac{x-vt-i0^+}{L}})^{-\alpha/4\kappa}, & t < 0 \end{cases} \\
 & = \frac{(L/2\pi)^{\alpha^2/2\kappa}}{[-i(x-vt)\text{sgn}(t) + 0^+]^{\alpha^2/4\kappa} [-i(-x-vt)\text{sgn}(t) + 0^+]^{\alpha/4\kappa}} \\
 & = \frac{(L/2\pi)^{2\gamma}}{(x^2 - v^2t^2 + i2vt\text{sgn}(t)0^+ + (0^+)^2)^\gamma} = \frac{(L/2\pi)^{2\gamma}}{(x^2 - v^2t^2 + i0^+)^\gamma}
 \end{aligned}$$

$\gamma = \lambda^2/4\kappa = \sqrt{V_1 V_2}/2V_1$ (choose the positive branch for $x \rightarrow \infty$).

- Imaginary-time ($\tau = it$) correlation is simplified $\frac{(L/2\pi)^{2\gamma}}{(z\bar{z})^\gamma}$, $z = x + iv\tau$

- 1d superfluid (boson condensation or $U(1)$ symm. breaking) only has an **algebraic long range order**, not real **long range order** (since $\langle : e^{i\theta(x,0)} :: e^{-i\theta(0,0)} : \rangle|_{x \rightarrow \infty} \not\rightarrow \text{const.}$) **Continuous symmetry cannot spontaneously broken in 1D. It can only “nearly broken”**

Correlation function and spectral function of $e^{i\theta} \sim \sigma^+$

$$\begin{aligned}
 G(x, t) &= i \langle T[: e^{i\theta(x,t)} :: e^{-i\theta(0,0)} :] \rangle \\
 &= i (1 - e^{i2\pi \frac{x-vt}{L} \text{sgn}(t)})^{-\gamma} (1 - e^{i2\pi \frac{-x-vt}{L} \text{sgn}(t)})^{-\gamma} \\
 &= \sum_n C_{m,n} i e^{i(m\frac{2\pi}{L}x - n\frac{2\pi v}{L}t) \text{sgn}(t)} = \sum_n C_{m,n} i e^{i(\kappa_m x - E_n t) \text{sgn}(t)} \\
 I(k, \omega) &= \sum_n C_{m,n} [\delta(k - \kappa_m) \delta(\omega - E_n) + \delta(k + \kappa_m) \delta(\omega + E_n)]
 \end{aligned}$$

Fourier transformation of $G(x, t)$:

$$\begin{aligned}
 &\int_0^L dx \int_{-\infty}^{\infty} dt e^{-i(kx - \omega t)} i e^{i(\kappa_m x - E_n t) \text{sgn}(t)} \\
 &= \int_0^L dx \int_0^{\infty} dt e^{-i[kx - (\omega + i0^+)t]} i e^{i(\kappa_m x - E_n t)} + (t < 0) \\
 &= \underbrace{\delta(k - \kappa_m)}_{L\delta_{k, \kappa_m}} \frac{i}{-i(\omega - E_n + i0^+)} = \underbrace{\delta(k - \kappa_m)}_{L\delta_{k, \kappa_m}} \left[\frac{-1}{\omega - E_n} + i\pi\delta(\omega - E_n) \right] \\
 I(k, \omega) &= \text{Im} G(k, \omega) / \pi
 \end{aligned}$$

Correlation function and spectral function of $e^{i\theta} \sim \sigma^+$

Correlation function of $e^{i\theta} \sim \sigma^+$

$$G(x,t) = \frac{i(L/2\pi)^{2\gamma}}{(x^2 - v^2 t^2 + i0^+)^{\gamma}} = \frac{i(L/2\pi)^{2\gamma}}{(y_1 y_2 + i0^+)^{\gamma}}$$

where $y_1 = x + vt$, $y_2 = x - vt$. We find

$$\begin{aligned} G(k, \omega) &= \int dx dt e^{-i(kx - \omega t)} \frac{i(L/2\pi)^{2\gamma}}{(x^2 - v^2 t^2 + i0^+)^{\gamma}} \\ &= \int dx dt e^{-i\frac{1}{2}[k(y_1 + y_2) - v^{-1}\omega(y_1 - y_2)]} \frac{i(L/2\pi)^{2\gamma}}{(y_1 y_2 + i0^+)^{\gamma}} \\ &\sim \int dy_1 dy_2 \frac{i e^{-i\frac{1}{2}[(k - \frac{\omega}{v})y_1 + (k + \frac{\omega}{v})y_2]}}{(y_1 y_2 + i0^+)^{\gamma}} \end{aligned}$$

up to a positive factor.

When taking the fractional power γ , choose the positive branch for $y_1 y_2 > 0$. For $y_1 y_2 < 0$, choose branch that connect to the positive branch for $y_1 y_2 > 0$. Now the term $i0^+$ becomes important.

Correlation function and spectral function of $e^{i\theta} \sim \sigma^+$

- $y_1 > 0, y_2 > 0$: Using $\int_0^\infty dx \frac{e^{-ax}}{x^\alpha} = \Gamma(1-\alpha)a^{\alpha-1}$, $\text{Re}(a) > 0$ and inserting 0^+ to make sure $\text{Re}(a) > 0$, we find

$$\begin{aligned} G_{++}(k, \omega) &= i \int_0^\infty dy_1 \int_0^\infty dy_2 \frac{e^{-i\frac{1}{2}(k-\frac{\omega}{v}-i0^+)y_1} e^{-i\frac{1}{2}(k+\frac{\omega}{v}-i0^+)y_2}}{(y_1 y_2 + i0^+)^\gamma} \\ &= i \left(\frac{i(k-\frac{\omega}{v}) + 0^+}{2} \right)^{\gamma-1} \Gamma(1-\gamma) \left(\frac{i(k+\frac{\omega}{v}) + 0^+}{2} \right)^{\gamma-1} \Gamma(1-\gamma) \\ &= i e^{i\frac{\pi}{2}(\gamma-1)[\text{sgn}(vk-\omega)+\text{sgn}(vk+\omega)]} \\ &\quad \left(\frac{|vk-\omega|}{2v} \right)^{\gamma-1} \left(\frac{|vk+\omega|}{2v} \right)^{\gamma-1} \Gamma^2(1-\gamma) \end{aligned}$$

Correlation function and spectral function of $e^{i\theta} \sim \sigma^+$

- $y_1 > 0, y_2 < 0$: Using $\int_0^\infty dx \frac{e^{-ax}}{x^\alpha} = \Gamma(1-\alpha)a^{\alpha-1}$, $\text{Re}(a) > 0$ and inserting 0^+ to make sure $\text{Re}(a) > 0$, we find

$$\begin{aligned}
 G_{+-}(k, \omega) &= i \int_0^\infty dy_1 \int_{-\infty}^0 dy_2 \frac{e^{-i\frac{1}{2}(k-\frac{\omega}{v}-i0^+)y_1} e^{-i\frac{1}{2}(k+\frac{\omega}{v}+i0^+)y_2}}{(y_1 y_2 + i0^+)^\gamma} \\
 &= i \int_0^\infty dy_1 \int_0^\infty dy_2 \frac{e^{-i\frac{1}{2}(k-\frac{\omega}{v}-i0^+)y_1} e^{i\frac{1}{2}(k+\frac{\omega}{v}+i0^+)y_2}}{(-y_1 y_2 + i0^+)^\gamma} \\
 &= i \left(\frac{i(k-\frac{\omega}{v})+0^+}{2} \right)^{\gamma-1} \left(\frac{-i(k+\frac{\omega}{v})+0^+}{2} \right)^{\gamma-1} e^{-i\pi\gamma} \Gamma^2(1-\gamma) \\
 &= i e^{-i\pi\gamma} e^{i\frac{\pi}{2}(\gamma-1)[\text{sgn}(vk-\omega)-\text{sgn}(vk+\omega)]} \\
 &\quad \left(\frac{|vk-\omega|}{2v} \right)^{\gamma-1} \left(\frac{|vk+\omega|}{2v} \right)^{\gamma-1} \Gamma^2(1-\gamma)
 \end{aligned}$$

Correlation function and spectral function of $e^{i\theta} \sim \sigma^+$

- $y_1 < 0, y_2 > 0$: Using $\int_0^\infty dx \frac{e^{-ax}}{x^\alpha} = \Gamma(1-\alpha)a^{\alpha-1}$, $\text{Re}(a) > 0$ and inserting 0^+ to make sure $\text{Re}(a) > 0$, we find

$$\begin{aligned}
 G_{-+}(k, \omega) &= i \int_{-\infty}^0 dy_1 \int_0^\infty dy_2 \frac{e^{-i\frac{1}{2}(k-\frac{\omega}{v}+i0^+)y_1} e^{-i\frac{1}{2}(k+\frac{\omega}{v}-i0^+)y_2}}{(y_1 y_2 + i0^+)^\gamma} \\
 &= i \int_0^\infty dy_1 \int_0^\infty dy_2 \frac{e^{i\frac{1}{2}(k-\frac{\omega}{v}+i0^+)y_1} e^{-i\frac{1}{2}(k+\frac{\omega}{v}-i0^+)y_2}}{(-y_1 y_2 + i0^+)^\gamma} \\
 &= i \left(\frac{-i(k-\frac{\omega}{v})+0^+}{2} \right)^{\gamma-1} \left(\frac{i(k+\frac{\omega}{v})+0^+}{2} \right)^{\gamma-1} e^{-i\pi\gamma} \Gamma^2(1-\gamma) \\
 &= i e^{-i\pi\gamma} e^{i\frac{\pi}{2}(\gamma-1)[- \text{sgn}(vk-\omega) + \text{sgn}(vk+\omega)]} \\
 &\quad \left(\frac{|vk-\omega|}{2v} \right)^{\gamma-1} \left(\frac{|vk+\omega|}{2v} \right)^{\gamma-1} \Gamma^2(1-\gamma)
 \end{aligned}$$

Correlation function and spectral function of $e^{i\theta} \sim \sigma^+$

- $y_1 < 0, y_2 < 0$: Using $\int_0^\infty dx \frac{e^{-ax}}{x^\alpha} = \Gamma(1-\alpha)a^{\alpha-1}$, $\text{Re}(a) > 0$ and inserting 0^+ to make sure $\text{Re}(a) > 0$, we find

$$\begin{aligned}
 G_{--}(k, \omega) &= i \int_{-\infty}^0 dy_1 \int_{-\infty}^0 dy_2 \frac{e^{-i\frac{1}{2}(k-\frac{\omega}{v}+i0^+)y_1} e^{-i\frac{1}{2}(k+\frac{\omega}{v}+i0^+)y_2}}{(y_1 y_2 + i0^+)^\gamma} \\
 &= i \int_0^\infty dy_1 \int_0^\infty dy_2 \frac{e^{i\frac{1}{2}(k-\frac{\omega}{v}+i0^+)y_1} e^{i\frac{1}{2}(k+\frac{\omega}{v}+i0^+)y_2}}{(y_1 y_2 + i0^+)^\gamma} \\
 &= i \left(\frac{-i(k-\frac{\omega}{v})+0^+}{2} \right)^{\gamma-1} \left(\frac{-i(k+\frac{\omega}{v})+0^+}{2} \right)^{\gamma-1} \Gamma^2(1-\gamma) \\
 &= i e^{i\frac{\pi}{2}(\gamma-1)[- \text{sgn}(vk-\omega) - \text{sgn}(vk+\omega)]} \\
 &\quad \left(\frac{|vk-\omega|}{2v} \right)^{\gamma-1} \left(\frac{|vk+\omega|}{2v} \right)^{\gamma-1} \Gamma^2(1-\gamma)
 \end{aligned}$$

Correlation function and spectral function of $e^{i\theta} \sim \sigma^+$

$$\begin{aligned}
 G(k, \omega) &\sim i \left(\frac{|vk - \omega|}{2v} \right)^{\gamma-1} \left(\frac{|vk + \omega|}{2v} \right)^{\gamma-1} \Gamma^2(1 - \gamma) \times \\
 &\left(e^{i\frac{\pi}{2}(\gamma-1)[\text{sgn}(vk-\omega)+\text{sgn}(vk+\omega)]} + e^{-i\pi\gamma} e^{i\frac{\pi}{2}(\gamma-1)[\text{sgn}(vk-\omega)-\text{sgn}(vk+\omega)]} \right. \\
 &\left. + e^{-i\pi\gamma} e^{i\frac{\pi}{2}(\gamma-1)[- \text{sgn}(vk-\omega)+\text{sgn}(vk+\omega)]} + e^{i\frac{\pi}{2}(\gamma-1)[- \text{sgn}(vk-\omega)-\text{sgn}(vk+\omega)]} \right) \\
 &= i \left(\frac{|vk - \omega|}{2v} \right)^{\gamma-1} \left(\frac{|vk + \omega|}{2v} \right)^{\gamma-1} \Gamma^2(1 - \gamma) \times
 \end{aligned}$$

$$\begin{cases}
 -e^{i\pi\gamma} + e^{-i\pi\gamma} + e^{-i\pi\gamma} - e^{-i\pi\gamma} = -2i \sin(\pi\gamma), & vk - \omega > 0, vk + \omega > 0 \\
 -e^{-i\pi\gamma} + e^{-i\pi\gamma} + e^{-i\pi\gamma} - e^{i\pi\gamma} = -2i \sin(\pi\gamma), & vk - \omega < 0, vk + \omega < 0 \\
 1 - 1 - e^{-i2\pi\gamma} + 1 = 1 - e^{-i2\pi\gamma}, & vk - \omega > 0, vk + \omega < 0 \\
 1 - e^{-i2\pi\gamma} - 1 + 1 = 1 - e^{-i2\pi\gamma}, & vk - \omega < 0, vk + \omega > 0
 \end{cases}$$

Spectral function:
$$I(k, \omega) = \left(\frac{|vk - \omega|}{2v} \right)^{\gamma-1} \left(\frac{|vk + \omega|}{2v} \right)^{\gamma-1} \Gamma^2(1 - \gamma) \times
 \begin{cases}
 0, & (\omega - vk)(\omega + vk) < 0 \\
 1 - \cos(2\pi\gamma), & (\omega - vk)(\omega + vk) > 0
 \end{cases}$$

$k = 0$ modes, and large momentum sectors

- Our theory so far contain only excitations described by oscillators a_k , $k = \frac{2\pi}{L} \times \text{int.}$.
- Our theory so far can produce excitation near $k = 0$, but not near $k = k_b = 2\pi \frac{N}{L}$.
- The correlation $\langle \mathcal{T} [: e^{i\theta(x,t)} :: e^{i\theta(0,0)} :] \rangle \sim (x^2 - v^2 t^2)^{-1/4\kappa} + 0 e^{ik_b x}$ contains nothing near k_b .
- To include the low energy sectors with large momentum, we need to include $k = 0$ modes:

Low energy excitations = ($k \neq 0$ modes) \otimes ($k = 0$ modes)

- Consider θ, φ non-linear σ -model:

$$L = \int dx \left(\frac{1}{2\pi} \partial_x \varphi + \frac{\bar{\phi}^2}{a} \right) \partial_t \theta - \frac{v}{4\pi} (\partial_x \theta)^2 - \frac{v}{4\pi} (\partial_x \varphi)^2$$

- The $k = 0$ sectors are labeled by $w_\theta, w_\varphi \in \mathbb{Z}$ (Only $q = \partial\varphi$ is physical):
 $\theta(x) = w_\theta \frac{2\pi}{L} x + \theta_0 + (k \neq 0 \text{ modes}), \quad \varphi(x) = w_\varphi \frac{2\pi}{L} x + (k \neq 0 \text{ modes}).$
 $L = (w_\varphi + \frac{\bar{\phi}^2 L}{a}) \dot{\theta}_0 - \frac{1}{2} \frac{2\pi}{L} v (w_\theta^2 + w_\varphi^2) \rightarrow E = \frac{1}{2} \frac{2\pi}{L} v (w_\theta^2 + w_\varphi^2)$



The physical meanings of winding numbers w_θ , w_φ from the connection to the lattice model

- **What is the meaning of w_φ (angular momentum of θ_0)?**

We note that $2\bar{\phi}a^{-1}q = \kappa\partial_x\varphi/\pi = \partial_x\varphi/2\pi = w_\varphi/L$.

So $w_\varphi = \int dx 2\bar{\phi}a^{-1}q = \sum_i 2\bar{\phi}q_i$

Spectral function of n_i

But what is $\sum_i 2\bar{\phi}q_i$? Remember that $\phi_i = \bar{\phi} + q_i$

and $|\phi_i\rangle = \frac{|\uparrow\rangle + \phi_i|\downarrow\rangle}{\sqrt{1+|\phi_i|^2}} = \frac{|0\rangle + \phi_i|1\rangle}{\sqrt{1+|\phi_i|^2}}$.

So $\langle n_i \rangle = \frac{|\phi_i|^2}{1+|\phi_i|^2} \approx |\phi_i|^2 \approx \bar{\phi}^2 + 2\bar{\phi}q_i$

Thus the canonical momentum of θ_0 ,


$\frac{\bar{\phi}^2 L}{a} + w_\varphi = \sum_i (\bar{\phi}^2 + 2\bar{\phi}q_i) = \sum_i n_i = N$, is the total

number of the bosons. This should be an exact result, since

$\theta_0 \sim \theta_0 + 2\pi$ and its angular momenta are quantized as integers.

- **What is the meaning of w_θ ?**

A non-zero w_θ gives rise $\phi_i = \bar{\phi}e^{iw_\theta x \frac{2\pi}{L}}$. Each boson carries momentum $w_\theta \frac{2\pi}{L}$. The total momentum is $w_\theta \frac{2\pi N_0}{L} = w_\theta k_b$.



Obtain the meanings of w_θ , w_φ within the field theory

$$L = \int dx \left(\frac{1}{2\pi} \partial_x \varphi + \frac{\bar{\phi}^2}{a} \right) \partial_t \theta - \frac{v}{4\pi} (\partial_x \theta)^2 - \frac{v}{4\pi} (\partial_x \varphi)^2$$

- The $U(1)$ symmetry transformation is given by $\theta \rightarrow \theta + \theta_0$. The angular momentum of θ_0 is the total number of the $U(1)$ charges (ie the number of bosons). From the corresponding Lagrangian $L = (w_\phi + \frac{\bar{\phi}^2 L}{a}) \dot{\theta}_0 + \dots$, we see the $U(1)$ charge is $Q = w_\phi - \frac{\bar{\phi}^2 L}{a}$
- The translation symmetry transformation is given by $\theta(x) \rightarrow \theta(x - x_0)$, $\varphi(x) \rightarrow \varphi(x - x_0)$. The canonical momentum of x_0 is the total momentum.
- We consider the field of form $\theta(x - x_0)$, $\varphi(x - x_0)$ and only x_0 is dynamical, ie time dependent (the $k \neq 0$ mode can be dropped):

$$\theta(x, t) = w_\theta \frac{2\pi}{L} (x + x_0(t)) + \theta_0 + (k \neq 0 \text{ modes}),$$


$$\varphi(x, t) = w_\varphi \frac{2\pi}{L} (x + x_0(t)) + (k \neq 0 \text{ modes}).$$

From the corresponding Lagrangian $L = (w_\phi + \frac{\bar{\phi}^2 L}{a}) \frac{2\pi}{L} w_\theta \dot{x}_0 + \dots$, we see the total momentum is $K = N \frac{2\pi}{L} w_\theta = k_b w_\theta$.

Winding-number changing operators

$$L = \int dx \left(\frac{1}{2\pi} \partial_x \varphi + \frac{\bar{\phi}^2 L}{a} \right) \partial_t \theta - \frac{v}{4\pi} (\partial_x \theta)^2 - \frac{v}{4\pi} (\partial_x \varphi)^2$$

- The **local operator** $e^{i\theta} = e^{i\alpha(\phi_1 + \phi_2)}$ changes the particle number N by -1 , i.e. change the winding number of φ , w_φ , by -1 .



- To see this explicitly

$$[\theta(x), \frac{1}{2\pi} \partial_y \varphi(y)] = i\delta(x - y)$$

We find $[\theta(x), \Delta\varphi] = i2\pi$ where $\Delta\varphi = \varphi(+\infty) - \varphi(-\infty)$.

Thus $\theta(x) = i2\pi \frac{d}{d\Delta\varphi} + \text{commutants of } \Delta\varphi$, and $e^{i\theta(x)} = e^{-2\pi \frac{d}{d\Delta\varphi} + \dots}$ is an operator that changes $\Delta\varphi$ by -2π , or w_φ by -1 , or particle number by -1

- Similarly, we have $[\theta(x), \varphi(y)] = -i2\pi\Theta(x - y)$

$$\rightarrow [\partial_x \theta(x), \varphi(y)] = -i2\pi\delta(x - y)$$

We find $[\Delta\theta, \varphi(y)] = -i2\pi$ where $\Delta\theta = \theta(+\infty) - \theta(-\infty)$.

Thus $\varphi(y) = i2\pi \frac{d}{d\Delta\theta}$, and $e^{i\varphi(x)} = e^{-2\pi \frac{d}{d\Delta\theta}}$ is an operator that changes $\Delta\theta$ by -2π , or change w_θ by -1 (i.e. total momentum by $-k_b$).

Local operators in 1D XY-model (superfluid)

- Lattice operators

$$\sigma_i^z = (\# \partial_x \theta + \# \partial_x \varphi) + \# e^{-i k_b x} e^{i \varphi(x)} + \dots$$

$$\sigma_i^+ = (\# + \# \partial_x \theta + \# \partial_x \varphi) e^{-i \theta(x)} + \# e^{-i k_b x} e^{-i \theta(x)} e^{i \varphi(x)} + \dots$$

- Set of local operators: $\partial_x \theta, \partial_x \varphi, \underbrace{e^{i(m_\theta \theta + m_\varphi \varphi)}}_{\text{change sectors}}$

or (from $\theta = \alpha(\phi_1 + \phi_2), \varphi = \beta(\phi_1 - \phi_2)$)

$$\partial_x \phi_1, \partial_x \phi_2, \underbrace{e^{i(m_1 \phi_1 + m_2 \phi_2)}}_{\text{change sectors}}$$

where $m_1 = \alpha m_\theta + \beta m_\varphi, m_2 = \alpha m_\theta - \beta m_\varphi$.

- **Fractionalization in XY-model (superfluid)**

A boson creation operator $\sigma^+ \sim e^{i\theta}$ (spin flip operator $\Delta S^z = 1$)

$$e^{i\theta} = e^{i\alpha(\phi_1 + \phi_2)}, \quad \phi_1 \text{ left-mover}, \quad \phi_2 \text{ right-mover}$$

$e^{i\alpha\phi_2}$ creates half boson (spin-1/2) in right-moving sector

$e^{i\alpha\phi_1}$ creates half boson (spin-1/2) in left-moving sector

Lattice translation and $U(1)$ symm. are not independent

- For a 1d superfluid of per-site-density n_b the ground state is described by a field $\phi(x) = \bar{\phi} e^{-i\theta(x)}$, $\theta(x) = 0$. The total momentum of the ground state is $K = 0$.
- We do a $U(1)$ symmetry twist: $\theta(L) = \theta(0) \rightarrow \theta(L) = \theta(0) + \Delta\theta$. The twisted state is described by a field $\theta(x) = \frac{\Delta\theta}{L}x$. The total momentum of the twisted state is $K = k_b \frac{\Delta\theta}{2\pi} = N + \Delta k = N \frac{\Delta\theta}{L}$.
- $U(1)$ symmetry twist = momentum boost $k_i \rightarrow k_i + \frac{\Delta\theta}{L}$.
Doing a symmetry twist operation in a symmetry can change the quantum number of another symmetry \rightarrow mixed anomaly
- A $2\pi U(1)$ symmetry twist can change the total crystal momentum by $k_b = 2\pi n_b$. Since 2π -crystal-momentum = 0-crystal-momentum, our bosonic system have an mixed translation- $U(1)$ anomaly when boson number per site $n_b \notin \mathbb{Z}$. \rightarrow **There is no translation and $U(1)$ symmetric product state.**
- We do a translation symmetry twist operation by adding ΔL sites \rightarrow change the total boson numbers (the $U(1)$ charges) of system by $n_b \Delta L$.


1d field theory – non-linear σ -model

- “Coordinate space” Lagrangian (rotor model): substitute one of the EOM $\frac{1}{2\pi} \partial_t \theta = \frac{1}{2\pi} V_2 \partial_x \varphi$ into the phase space Lagrangian

$$L = \int dx \frac{V_2^{-1}}{4\pi} (\partial_t \theta)^2 - \frac{V_1}{4\pi} (\partial_x \theta)^2 + \underbrace{\frac{\bar{\phi}^2}{a} \partial_t \theta}_{\text{a topo. term}}$$

$$= \int dx \frac{V_2^{-1}}{4\pi} (iu^* \partial_t u)^2 - \frac{V_1}{4\pi} (iu^* \partial_x u)^2 - i \frac{\bar{\phi}^2}{a} u^* \partial_t u$$

- The field is really $u = e^{i\theta}$, not θ . The above is the so called non-linear σ -model, where the field is a map from space-time manifold to the **target space** S^1 : $M_{\text{space-time}}^{d+1} \rightarrow U(1)$.
- In general, the target space is the symmetric space $G_{\text{symm}}/G_{\text{unbroken}}$ (the minima of the symmetry breaking potential).
- The topological term $i \frac{\bar{\phi}^2}{a} u^* \partial_t u$ cannot be dropped (since it is not a total derivative). When $\bar{\phi}^2 = n \notin \mathbb{Z}$, the topological term makes it impossible for the non-linear σ -model to have a gapped phase (an effect of mixed anomaly between $U(1)$ symmetry and translation symmetry).
- The above is a **low energy effective theory** for $U(1)$ symm breaking



© CPH Theory. All rights reserved. This content is excluded from our Creative Commons license. For more information, see <https://ocw.mit.edu/help/faq-fair-use>.

Symmetry, gauging, and conservation

- Consider a system described by a complex field u

$$S = \int dt dx \mathcal{L}(u)$$

with $U(1)$ symmetry: $\mathcal{L}(e^{i\lambda}u) = \mathcal{L}(u)$. We like to show that the system has a conserved current j^μ , $\mu = t, x$: $\partial_t j^t + \partial_x j^x = \partial_\mu j^\mu = 0$.

- **Gauge the $U(1)$ symmetry:**

- $u(x) \rightarrow e^{i\lambda_I(x)}u(x)$ gives rise to $u_I^* \partial_\mu u_I \rightarrow u_I^* (\partial_\mu + i\partial_\mu \lambda_I) u_I$, $\mu = t, x$.
- Replacing $\partial_\mu \lambda_I$ by a vector potential A_μ^I : $u_I^* (\partial_\mu + iA_\mu^I) u_I$ gives rise to a gauged theory $\mathcal{L} \rightarrow \mathcal{L}(u, A_\mu)$. Here A_μ is viewed as non-dynamical background field. We have

$$\mathcal{L}(u, A_\mu) = \mathcal{L}(e^{i\lambda}u, A_\mu - \partial_\mu \lambda)$$

- The $U(1)$ current of the gauged theory (setting $A_\mu = 0$ gives rise to the $U(1)$ current of the original theory)

$$\delta S = \int dt dx j^\mu \delta A_\mu, \quad j^\mu = \frac{\delta \mathcal{L}(u, A_\mu)}{\delta A_\mu}.$$

Symmetry, gauging, and conservation

- The current conservation:

$$\begin{aligned}\delta S &= \int d^2x^\mu \mathcal{L}(e^{i\lambda}u, A_\mu) - \mathcal{L}(u, A_\mu) \\ &= \int d^2x^\mu \mathcal{L}(u, A_\mu + \partial_\mu\lambda) - \mathcal{L}(u, A_\mu) = \int d^2x^\mu j^\mu \partial_\mu\lambda = - \int d^2x^\mu \lambda \partial_\mu j^\mu\end{aligned}$$

If $u(x, t)$ satisfies the equation of motion, then the corresponding $\delta S = 0$. This allows us to show the existence of a conserved current

$$\partial_\mu j^\mu(u) = 0.$$

- **Example:** $\partial_\mu\theta = -iu^*\partial_\mu u \rightarrow \partial_\mu\theta + A_\mu = -iu^*(\partial_\mu + iA_\mu)u$

$$\mathcal{L} = \frac{V_2^{-1}}{4\pi}(\partial_t\theta)^2 - \frac{V_1}{4\pi}(\partial_x\theta)^2 + \frac{\bar{\phi}^2}{a}\partial_t\theta$$

$$\rightarrow \mathcal{L} = \frac{V_2^{-1}}{4\pi}(\partial_t\theta + A_t)^2 - \frac{V_1}{4\pi}(\partial_x\theta + A_x)^2 + \frac{\bar{\phi}^2}{a}(\partial_t\theta + A_t)$$

$$\rightarrow j^\mu = \frac{\delta\mathcal{L}}{\delta A_\mu}, \quad j^t = \frac{V_2^{-1}}{2\pi}(\partial_t\theta + A_t), \quad j^x = -\frac{V_1}{2\pi}(\partial_x\theta + A_x).$$

Another example of gauging symmetry

Consider the following effective theory for 1d bosonic superfluid

$$\begin{aligned} L &= \int dx \frac{K_{IJ}}{4\pi} \partial_x \varphi_I \partial_t \varphi_J - \frac{V_{IJ}}{4\pi} \partial_x \varphi_I \partial_x \varphi_J + q_I \partial \phi_I \\ &= \int dx \frac{K_{IJ}}{4\pi} \partial_x u_I^* \partial_t u_J - \frac{V_{IJ}}{4\pi} \partial_x u_I^* \partial_x u_J - i q_I u_I^* \partial_t u_I \end{aligned}$$

$$I, J = 1, 2, \quad \varphi_I \sim \varphi_I + 2\pi, \quad u_I = e^{i\varphi_I}, \quad K = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad q = \begin{pmatrix} \frac{\bar{\phi}^2}{a} \\ 0 \end{pmatrix}.$$

- The effective field theory has two $U(1)$ symmetries:
 - $\varphi_1 \rightarrow \varphi_1 + \lambda_1$ for boson number conservation
Conjugate of λ_1 is $\int dx \frac{1}{2\pi} \partial_x \varphi_2 = w_\varphi = N$.
 - $\varphi_2 \rightarrow \varphi_2 + \lambda_2$ for momentum conservation.
Conjugate of λ_2 is $\int dx \frac{1}{2\pi} \partial_x \varphi_1 = w_\theta = K/k_b$.

Another example of gauging symmetry

- Gauging the two $U(1)$ symmetries:
 - $u_I(x) \rightarrow e^{i\lambda_I(x)} u_I(x)$ gives rise to $u_I^* \partial_\mu u_I \rightarrow u_I^* (\partial_\mu + i\partial_\mu \lambda_I) u_I$, $\mu = t, x$.
 - Replacing $\partial_\mu \lambda_I$ by a vector potential A_μ^I gives rise to a gauged theory

$$\begin{aligned}\mathcal{L} &= \frac{K_{IJ}}{4\pi} (\partial_x - iA_x^I) u_I^* (\partial_t + iA_t^J) u_J - \frac{V_{IJ}}{4\pi} (\partial_x - iA_x^I) u_I^* (\partial_x + iA_x^J) u_J \\ &\quad - iq_I u_I^* (\partial_t + iA_t^I) u_I \\ &= \frac{K_{IJ}}{4\pi} (\partial_x \varphi_I + A_x^I) (\partial_t \varphi_J + A_t^J) - \frac{V_{IJ}}{4\pi} (\partial_x \varphi_I + A_x^I) (\partial_x \varphi_J + A_x^J) + q_I (\partial_t \varphi_I + A_t^I)\end{aligned}$$

- Conserved current

$$j_I^t = \frac{K_{IJ}}{4\pi} (\partial_x \varphi_J + A_x^J) + q_I, \quad j_I^x = \frac{K_{IJ}}{4\pi} (\partial_t \varphi_J + A_t^J) - \frac{V_{IJ}}{2\pi} (\partial_x \varphi_J + A_x^J)$$

- Equation of motion \rightarrow conservation

$$\begin{aligned}& - \frac{K_{IJ}}{4\pi} \partial_x (\partial_t \varphi_J + A_t^J) - \frac{K_{IJ}}{4\pi} \partial_t (\partial_x \varphi_J + A_x^J) + \frac{V_{IJ}}{2\pi} \partial_x (\partial_x \varphi_J + A_x^J) = 0 \\ \rightarrow & - \partial_t j_I^t - \partial_x j_I^x = 0\end{aligned}$$

Symmetry twist, pumping, and anomaly

- But for certain background field $A'_\mu(x, t)$, the equation of motion cannot be satisfied \rightarrow non-conservation. **Symmetry twist** \rightarrow **Pumping**
Background field $A'_\mu(x, t) =$ symmetry twist.
Non-conservation = pumping
- Consider $A'_t = 0$, A'_x independent of x , but dependent on t . Equation of motion becomes

$$-\frac{K_{IJ}}{2\pi} \partial_x \partial_t \varphi_J + \frac{V_{IJ}}{2\pi} \partial_x^2 \varphi_J = \frac{K_{IJ}}{4\pi} \partial_t A'_x$$

It has no solution since, on a ring of size L ,

$$0 = \int_0^L dx \left[-\frac{K_{IJ}}{2\pi} \partial_x \partial_t \varphi_J + \frac{V_{IJ}}{2\pi} \partial_x^2 \varphi_J \right] = \int_0^L dx \frac{K_{IJ}}{4\pi} \partial_t A'_x \neq 0$$

- The non-zero pumped $U(1)$ charge $\rightarrow U(1)$ anomaly

$$\dot{Q}_I = \int_0^L dx \partial_t j_I^t = \int_0^L dx \partial_t \left[\frac{K_{IJ}}{4\pi} (\partial_x \varphi_J + A'_x) + q_I \right] = \int_0^L dx \partial_t \frac{K_{IJ}}{4\pi} A'_x$$

Anomaly and mixed anomaly

Consider chiral boson theory

$$L = \int dx \frac{K_{IJ}}{4\pi} \partial_x \varphi_I \partial_t \varphi_J - \frac{V_{IJ}}{4\pi} \partial_x \varphi_I \partial_x \varphi_J + q_I \partial \phi_I$$

$$\dot{Q}_I = \int_0^L dx \partial_t j_I^t = \int_0^L dx \partial_t \frac{K_{IJ}}{4\pi} A_x^J$$

- $K = (1)$, the theory is actually fermionic and describes a chiral fermion.
 - The $U(1)$ symmetry twist pumps the $U(1)$ charge $\rightarrow U(1)$ anomaly
- $K = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$, the theory is non-chiral describing 1d bosonic superfluid.
 - The first $U(1)$ symmetry twist does not pump the first $U(1)$ charge. The first $U(1)$ is not anomalous.
 - The second $U(1)$ symmetry twist does not pump the second $U(1)$ charge. The second $U(1)$ is not anomalous.
 - The first $U(1)$ symmetry twist pumps the second $U(1)$ charge. The $U(1) \times U(1)$ symmetry has a mixed anomaly.

Anomaly and mixed anomaly

- $K = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$, the theory is non-chiral describing 1d Fermi liquid.
 - The first $U(1)$ symmetry twist pumps the first $U(1)$ charge. The first $U(1)$ is anomalous.
 - The second $U(1)$ symmetry twist pumps the second $U(1)$ charge. The second $U(1)$ is anomalous.
- The “+” $U(1)$: $\varphi_1 \rightarrow \varphi_1 + \lambda_+$, $\varphi_2 \rightarrow \varphi_2 + \lambda_+$ → the fermion number
The “-” $U(1)$: $\varphi_1 \rightarrow \varphi_1 + \lambda_-$, $\varphi_2 \rightarrow \varphi_2 - \lambda_-$ → the total momentum
provided that the fermion density is not zero.
- The “+” $U(1)$ symmetry twist does not pump the “+” $U(1)$ charge. The “+” $U(1)$ is not anomalous.
 - The “-” $U(1)$ symmetry twist does not pump the “-” $U(1)$ charge. The “-” $U(1)$ is not anomalous.
 - The “+” $U(1)$ symmetry twist does not pump the “-” $U(1)$ charge. There is a mixed anomaly between “+” $U(1)$ and “-” $U(1)$ symmetries. **The $U^2(1)$ symmetric state must be gapless.**

Why $K = (1)$ chiral boson theory describes chiral fermions

$K = (1)$ chiral boson field theory:

$$\begin{aligned} L &= \int dx \frac{1}{4\pi} \partial_x \varphi \partial_t \varphi - \frac{V}{4\pi} \partial_x \varphi \partial_x \varphi \\ &= \sum_{k=-\infty}^{+\infty} \frac{-i}{4\pi} k \varphi_{-k} \dot{\varphi}_k - \frac{V}{4\pi} k^2 \varphi_{-k} \varphi_k, \quad \varphi(x) = \sum_{k=-\infty}^{+\infty} \frac{e^{ikx}}{\sqrt{L}} \varphi_k \\ &= \sum_{k>0} \frac{-i}{2\pi} k \varphi_{-k} \dot{\varphi}_k - \frac{V}{2\pi} k^2 \varphi_{-k} \varphi_k \end{aligned}$$

The canonical conjugate of φ is $\frac{1}{4\pi} \partial_y \varphi(y)$ or $\frac{1}{2\pi} \partial_y \varphi(y)$

$$[\varphi_k, \frac{-ik'}{2\pi} \varphi_{-k'}] = i \delta_{k-k'},$$

$$[\varphi(x), \frac{1}{2\pi} \partial_y \varphi(y)] = i \sum_k L^{-1} e^{ik(x-y)} = i \int \frac{dk}{2\pi} e^{ik(x-y)}$$

$$[\varphi(x), \frac{1}{2\pi} \partial_y \varphi(y)] = i \delta(x-y), \quad [\varphi(x), \varphi(y)] = i\pi \text{sgn}(x-y).$$

Why $K = (1)$ chiral boson theory describes chiral fermions

- $\varphi(x)$ is a compact field $\varphi(x) \sim \varphi(x) + 2\pi$. Thus $\varphi(x)$ is not an allowed operator. $e^{\pm i\varphi(x)}$ are allowed operators, all other allowed operators are generated by $e^{\pm i\varphi(x)}$.
- The allowed operators are non-local and should be forbidden:

$$\begin{aligned} e^{i\varphi(x)} e^{i\varphi(y)} &= e^{[i\varphi(x), i\varphi(y)]} e^{i\varphi(y)} e^{i\varphi(x)} \\ &= e^{i\pi \text{sgn}(x-y)} e^{i\varphi(y)} e^{i\varphi(x)} = -e^{i\varphi(y)} e^{i\varphi(x)} \end{aligned}$$

- Or we regard the non-local operators $e^{\pm i\varphi(x)}$ as local fermion operator, and regard the chiral boson theory as a theory for fermions.
- The imaginary-time (time-ordered) correlation function for $e^{\pm i\varphi(x)}$:

$$\langle e^{-i\varphi(x,\tau)} e^{i\varphi(0)} \rangle \sim \frac{1}{x + i\tau} = \frac{1}{z}$$

which is identical to the correlation function of free chiral fermion $c(x, t)$, and allows us to identify $c(x, t) \sim e^{i\varphi(x, t)}$.

$K = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ boson theory describes 1d Fermi liquid

- **Bosonization:**

$$L = \int dx \frac{1}{4\pi} \partial_x \varphi_R \partial_t \varphi_R - \frac{v_F}{4\pi} \partial_x \varphi_R \partial_x \varphi_R - \frac{1}{4\pi} \partial_x \varphi_L \partial_t \varphi_L - \frac{v_F}{4\pi} \partial_x \varphi_L \partial_x \varphi_L + q \partial_t (\varphi_R + \varphi_L)$$

describes 1d non-interacting fermions with Fermi velocity k_F .

- The fermion number $U(1)$ symmetry: $\varphi_R \rightarrow \varphi_R + \theta$, $\varphi_L \rightarrow \varphi_L + \theta$.

The canonical conjugate of θ is the fermion number \rightarrow Fermion number density is given by $n_F = \frac{1}{2\pi} (\partial_x \varphi_R - \partial_x \varphi_L)$.

- Interacting 1d fermions via bosonization:

$$L = \int dx \frac{1}{4\pi} \partial_x \varphi_R \partial_t \varphi_R - \frac{v_F}{4\pi} \partial_x \varphi_R \partial_x \varphi_R - \frac{1}{4\pi} \partial_x \varphi_L \partial_t \varphi_L - \frac{v_F}{4\pi} \partial_x \varphi_L \partial_x \varphi_L + \frac{V}{(2\pi)^2} (\partial_x \varphi_R - \partial_x \varphi_L)^2 + q \partial_t (\varphi_R + \varphi_L)$$

describes 1d interacting fermions, which allow us to compute fermion correlation $\langle c(x, t) c^\dagger(0) \rangle$, etc .

Fractionalization in general 1d chiral boson theory

$$L = \int dx \frac{K_{IJ}}{4\pi} \partial_x \varphi_I \partial_t \varphi_J - \frac{V_{IJ}}{4\pi} \partial_x \varphi_I \partial_x \varphi_J, \quad \varphi_I \sim \varphi_I + 2\pi,$$

with K_{II} even. The canonical conjugate of φ_I is $\frac{K_{IJ}}{2\pi} \partial_x \varphi_J \rightarrow$

$$[\varphi_I(x), \varphi_J(y)] = i\pi (K^{-1})_{IJ} \text{sgn}(x - y)$$

- All the allowed operators have the form $e^{i l_I \varphi_I(x)}$ where $l_I \in \mathbb{Z}$. The commutation of allowed operators

$$e^{i l_I \varphi_I(x)} e^{i \tilde{l}_J \varphi_J(y)} = e^{i \pi \tilde{l} K^{-1} l} e^{i \tilde{l}_J \varphi_J(y)} e^{i l_I \varphi_I(x)}$$

- Moving operator $e^{i l_I \varphi_I(x)}$ around $e^{i \tilde{l}_J \varphi_J(y)}$ induce a phase $e^{i 2\pi \tilde{l} K^{-1} l} \rightarrow$ **mutual statistics**. The imaginary-time correlation between $e^{i l_I \varphi_I(x)}$ and $e^{i \tilde{l}_J \varphi_J(y)}$ has a form

$$\langle \dots e^{i l_I \varphi_I(z_1)} e^{i \tilde{l}_J \varphi_J(z_2)} \dots \rangle \sim \frac{1}{(z_1 - z_2)^\gamma (\bar{z}_1 - \bar{z}_2)^{\bar{\gamma}}}, \quad \gamma - \bar{\gamma} = \tilde{l} K^{-1} l.$$

Fractionalization in general 1d chiral boson theory

Most of the allowed operators $e^{iI\varphi_I(x)}$ are not local (ie far away operators do not commute)

- **Local operators**: the operators $e^{iI^{loc}\varphi_I(x)}$ that commute with all allowed operator that are far way:

$$I^{loc}K^{-1}I = \text{even int.} \quad \forall I \in \mathbb{Z} \quad \rightarrow \quad I^{loc} = K_{IJ}n_J.$$

$e^{iI^{loc}\varphi_I(x)}$ corresponds to lattice boson operators.

- The allowed non-local operator $e^{iI\varphi_I(x)}$ create quasi particle with fractional statistics given by $e^{i\pi IK^{-1}I}$.
- In fact, the chiral boson model for most K is anomalous, ie can not be realized by 1d lattice boson model. But it can be realized by the boundary of 2d FQH Hall state. So the chiral boson model is a edge theory of 2d 2d FQH Hall state.

MIT OpenCourseWare

<https://ocw.mit.edu>

8.513 Modern Quantum Many-body Physics for Condensed Matter Systems

Fall 2021

For information about citing these materials or our Terms of Use, visit:

<https://ocw.mit.edu/terms>.